






Source: EEA and UNEP, Priority issues in the Mediterranean environment, 2006.



Coastal pollution of the Mediterranean



(UNEP/MAP, 2012; State of the Mediterranean Marine and Coastal Environment)























Hydrogen 1

The Russian chemist, Dmitri Mendeleyev, was the first to observe that if elements were listed in order of atomic mass, they showed regular (periodical) repeating properties. He formulated his discovery in a periodic table of elements, now regarded as the backbone of modern chemistry.

The crowning achievement of Mendeleyev's periodic table lay in his prophecy of then, undiscovere elements. In 1869, the year he published his periodic classification, the elements gallium, germanium and scandium were unknown. Mendeleyev left spaces for them in his table and even predicted their atomic masses and other chemical properties. Six years later, gallium was discovered and his predictions were found to be accurate. Other discoveries followed and their chemical behaviour matched that predicted by Mendeleyev.

This remarkable man, the youngest in a family of 17 children, has left the scientific community with a classification system so powerful that it became the cornerstone in chemistry teaching and the prediction of new elements ever since. In 1955, element 101 was named after him: Md, Mendelevium.





















































































































































































































Geographical distribution of *Posidonia oceanica* (Michel, 2012; redrawn using data from Lipkin et al., 2003; Procaccini et al., 2003; Boudouresque et al., 2006; Gobert et al., 2006; Meinesz et al., 2009). 1: Gibraltar; 2: Almeria; 3: Oran; 4: Coasts of Syria, Israel and Lebanon; A: Rhone estuary; B: Po estuary; C: Nile estuary.





Objectives

- i. to assess, for the first time, the TE contamination in the entire Mediterranean biomonitored with *P. oceanica*;
- ii. to propose a new 5-level water quality scale;
- iii. to calculate coastal water pollution indices from bioaccumulated TE levels measured in that bioindicator;
- iv. to produce, on the basis of these data, accurate maps of the contamination of the Mediterranean by TEs.

Overall aim of this work:


to develop a holistic approach in order to provide scientists, stakeholders and decision makers with a global tool to bioassess the TE contamination severity of Mediterranean coastal waters.





Sampling

110 sites differing by their levels of exposure to contaminants sampled for *P. oceanica*.












P. oceanica compartmentalization



Left: *P. oceanica* shoots fixed on a plagiotropic rhizome. Right: (A) shoot of leaves on a plagiotropic rhizome; (B, C) adult leaves; (D) intermediate leaf; (E) juvenile leaf (modified after Libes and Boudouresque, 1987).

Ε

T

Н

0


D

S





Trace elements monitored



















E

Н

0

D

S





Maths and stats: monitoring

- Water quality scale
- Contamination indices:
 - Trace Element Spatial Variation Index (TESVI)
 - Trace Element Pollution Index (TEPI)

Maths and stats: spatial analysis

- GIS mapping
- Principal component analysis
- Cluster analysis
- Correlation analysis



Maths and stats: 5-level water quality scale



qu. 1	qu. 2	qu. 3	qu. 4	
0.6000	1.3333	1.7635	2.9333	
0.6333	1.3333	1.7667	2.9667	
0.6667	1.3667	1.7682	3.0012	
0.7000	1.4000	1.8000	3.1253	
0.7667	1.4000	1.8000	3.2113	
0.8000	1.4000	1.8018	3.2333	
0.8333	1.4333	1.8241	3.2333	
0.8333	1.4333	1.8349	3.3667	
0.8333	1.4333	1.8667	3.3667	
0.8667	1.4511	1.9667	3.4705	
0.9000	1.4667	1.9667	3.6000	
1.0333	1.5000	2.0000	3.7667	
1.0667	1.5000	2.0333	3.8000	
1.0667	1.5180	2.1667	3.9000	
1.1333	1.5667	2.2000	4.0667	
1.1667	1.6000	2.2793	4.0667	
1.1667	1.6000	2.3333	4.4333	
1.1667	1.6333	2.3333	4.6000	
1.2000	1.6333	2.3667	5.3333	
1.2333	1.6333	2.4000	5.4790	
1.2333	1.6667	2.4333	5.5667	
1.2667	1.6667	2.5333	5.9129	
1.2667	1.7000	2.6333	5.9250	
1.2667	1.7000	2.6667	6.0751	
1.2667	1.7000	2.7333	6.1230	
1.2667	1.7333	2.9333	7.9000	
1.3000	1.7333	2.9333	8.5667	
1.3000			14.5000	

Quartile means

Superior	limit	of
quartiles	Pb	
quartile 1	1.308	33
quartile 2	1.748	34
quartile 3	2.933	33
quartile 4	14.500	00


5 contamination levels

< 1st qu. mean : very low CL	
1 st -2 nd qu. mean: low CL	
2 nd -3 rd qu. mean: medium CL	
3 rd -4 th qu. mean: high CL	
> 4 th qu. mean: very high CL	




UMONS STARESO STARESO A R C A P M E D


Sub-areas differences




















Maths and stats: pollution indices



























Maths and stats: pollution indices













- ❖ How to order and to compare TEs according to the overall spatial variability of their environmental levels through the whole of a studied area?
- How to compare global pollution levels in TEs between several monitored sites?














Sampling

Pollution indices developed first for the French Mediterranean littoral.



Marine Pollution Bulletin 89 (2014) 390-406



Contents lists available at ScienceDirect


Marine Pollution Bulletin

journal homepage: www.elsevier.com/locate/marpolbul

A reassessment of the use of *Posidonia oceanica* and *Mytilus galloprovincialis* to biomonitor the coastal pollution of trace elements: New tools and tips

J. Richir*, S. Gobert

Laboratory of Oceanology, MARE Centre, University of LIEGE, B6C, 4000 LIEGE, Sart Tilman, Belgium





Α



Trace Element Spatial Variation Index

How to order and to compare TEs according to the overall spatial variability of their environmental levels through the whole of a studied area?

Trace Element Spatial Variation Index (TESVI)

$$TESVI = [(x_{max}/x_{min}) / (\sum (x_{max}/x_{i})/n)] * SD$$

where:

- x_{max} and x_{min} are the maximum and minimum mean concentrations recorded among the n sites,
- x_i are mean concentrations recorded in each of the n sites,
- SD is the standard deviation of the weighted sum $\sum (x_{max}/x_i)/n$.

The highest the index value, the more environmental levels of a TE globally vary (punctual contaminations and overall coastal spatial heterogeneity of TE levels taken into account) through the whole of the studied area the index is applied to.






TESVI value calculation

Trace Element Spatial Variation Index (TESVI)

$$TESVI = [(x_{max}/x_{min}) / (\sum (x_{max}/x_i)/n)] * SD$$








TESVI value calculation

Trace Element Spatial Variation Index (TESVI)

$$TESVI = [(x_{max}/x_{min}) / (\sum (x_{max}/x_i)/n)] * SD$$







TESVI graphical representation

A. TEs broadly monitored with P. oceanica


TES broadly monitored with 7. oceanica					
	x_{max}/x_{min}	$\sum (x_{max}/x_i)/18 \pm SD$ TESVI Site x_n		Site x _{max}	
Cr	6.0	3.6 ± 1.3	2.2	St Florent	
Fe	4.4	2.0 ± 0.9	1.9	Bravone	
Ni	2.4	1.6 ± 0.3	(0),(5)	St Raphaël	
Cu	3.4	1.9 ± 0.7	1.2	Villefranche	
Zn	19.6	13.3 ± 4.4	6.5	Bravone	
Cd	3.9	1.9 ± 0.7	1.4	St Raphaël	
Pö	4.4	2.7 ± 1.2	2,0	Ajaccio N.	
Be	3.1	1.6 ± 0.6	1.0	Ajaccio N.	
Al	7.5	2.2 ± 1.8	6.1	Ajaccio N.	
W	14.5	5.9 ± 5.0	12.3	Antibes	
Mn	2.2	1.6 ± 0.4	0.5	St Raphaël	
Co	2.9	1.8 ± 0.5	0.7	St Raphaël	
As	10.6		4.9	P. des Chèvres	
Se	1.7		0.3	Calvi	
Mo	22.8	22.8	10.5	Aregno	
Ag	3.1	(EE.U)	0.9	La Vesse	
Sn (BAL)	6.9		3.8	Corbière	
Sb	4.4	3.6 ± 0.7	0.9	Bravone	
Bi	13.6	6.1 ± 3.5	7.9	P. des Chèvres	

TESVI values were listed in ascending order:

Se, Ni, Mn, Co, Sb, Ag, Be, Cu, Cd, Fe, Pb, Cr, Sn, As, Al, Zn, Bi, Mo, V



TESVI graphical representation CAPMED




TESVI values were listed in ascending order: Se, Ni, Mn, Co, Sb, Ag, Be, Cu, Cd, Fe, Pb, Cr, Sn, As, Al, Zn, Bi, Mo, V





TESVI graphical representation




TESVI values were listed in ascending order: Se, Ni, Mn, Co, Sb, Ag, Be, Cu, Cd, Fe, Pb, Cr, Sn, As, Al, Zn, Bi, Mo, V





Sampling

110 sites differing by their levels of exposure to contaminants sampled for *P. oceanica*.





D


U

S

S


О N


Spatial variability





TE	TESVI
Hg	3.9
Cd	8.7
Cu	9.2
Pb	13.3
As	29.4
Ag	34.9
Ni	92.7


Proportional ordinate **scaling** between TEs:





















Trace Element Pollution Index

How to compare global pollution levels in TEs between several monitored sites?

Trace Element Pollution Index (TEPI)

TEPI =
$$(Cf_1 * Cf_2 ... Cf_n)^{1/n}$$
,

a weighted version of the Metal Pollution Index (MPI) of Usero et al. (1996, MPB)

where:


• Cf_n is the mean normalized concentration of the TE n in a given monitored site.

The highest the index value, the more the monitored site is globally contaminated in TEs compared to the others.



Mean normalization







Random list 1:

Be, Al, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Se, Mo, Ag, Cd, Sn, Sb, Pb, Bi

P. Chèvres (4)







Random list 5:

Sb, V, Sn, Se, Be, Pb, Ni, Mo, Mn, Fe, Cu, Cr, Co, Cd, Bi, Zn, As, Ag, Al

P. Chèvres (4)











Sampling

110 sites differing by their levels of exposure to contaminants sampled for *P. oceanica*.




Global contamination





Sites





0

D

S





Maths and stats: monitoring

- Water quality scale
- Contamination indices:
 - Trace Element Spatial Variation Index (TESVI)
 - Trace Element Pollution Index (TEPI)


Maths and stats: spatial analysis

- E T
 - GIS mapping
 - Principal component analysis
 - Cluster analysis
 - Correlation analysis





Principal component analysis








Correlation analysis

	Ag	As	Cd	Cu	Hg	Ni	Pb	TEPI
Ag	1.000							
As	0.204	1.000						
Cd	0.291	-0.168	1.000					
Cu	0.549	0.096	0.084	1.000				
Hg	0.008	-0.074	-0.043	-0.047	1.000			
Ni	0.464	0.178	0.625	0.425	0.006	1.000		
Pb	0.294	0.252	0.110	0.261	-0.014	0.126	1.000	
TEPI	0.777	0.512	0.383	0.539	0.132	0.686	0.518	1.000

Correlation matrix of parametric Pearson's correlation coefficients and non-parametric Spearman's rank correlation coefficients between TE concentrations and TEPI values in P. oceanica. Correlations significant at p < 0.05 are in bold; correlations significant at p < 0.01 are in bold underlined.




PRODUCED BY THE FOUNDATION FOR EDUCATION, SCIENCE AND TECHNOLOGY FOR NATIONAL SET WEEK 200





Cluster analysis



Dendrographic classification after cluster analysis of Ag, As, Cd, Cu, Hg, Ni and Pb concentrations measured in the blades of *Posidonia oceanica*, and TEPI values calculated from mean normalized TE concentrations.


R E S U D U S S 0

N






Cluster analysis




TEPI values calculated from mean normalized Ag, As, Cd, Cu, Hg, Ni and Pb concentrations measured in the blades of *P. oceanica* adult leaves. Sampling sites are sorted on the graph according to the dendrographic classification after cluster analysis of trace element concentrations and TEPI values.














Conclusion

Sustainable coastal management requires the development of appropriate contamination classification systems.



- water quality scale;
- pollution index (TEPI and TESVI);
- spatial analysis (PCA, CA, correlation analysis and GIS mapping).

In the Mediterranean:

- To assess TE contamination threats;
- To depict contamination gradients;
- To monitor TE contamination at regional or local scales.

<u>Conclusion</u>: to privilege such <u>holistic approaches</u> to accurately monitor the TE contamination rate of coastal waters and to transfer <u>relevant information</u> on this <u>composite problem</u> to environmental managers and policy makers.





Acknowledgements

This study was supported by the collaboration of several laboratories from all around the Mediterranean:

- Pr Gérard Pergent, Pr Céline Lafabrie, Dr Christine Pergent-Martini, Dr Maylis Salivas-Decaux and Dr Cecilia Lopez y Royo, University of Corte (Corsica, France);
- Dr Saïd Belbacha and Dr Rachid Semroud (Algeria);
- University of Zagreb, Dr Tatjana Bakran-Petricioli (Croatia);
- Melina Marcou and Myroula Hadjichristoforou (Cyprus);
- Dr Stéphane Sartoretto, Ifremer Center of Toulon,
- Berangère Casalta (France);
- Dr Georges Skoufas and Dr Eugenia Apostolaki, Institute of Oceanography, Hellenic Centre for Marine Research (Greece);
- Dr Cristina Buia and Dr Pier Panzalis (Italy);
- Dr Joseph Borg (Malta);
- Robert Turk, Borut Mavric and Lovrenc Lipej (Slovenia);
- Boris Weitzmann, Dr Yolanda Fernandez Torquemade, F. Javier Zapata Salgado, Elvira Alvarez Pérez, Dr Javier Romero, Izaskun Llagostera (Spain);
- Dr Habib Langar (Tunisia); Dr Baris Akçalı and Billür Celebi (Turkey). We also thank Steven Barillec, Nicolas Luy and Renzo Biondo for their help.



More info ...



Journal of Environmental Management 151 (2015) 486-499



Contents lists available at ScienceDirect

Journal of Environmental Management





Bioassessment of trace element contamination of Mediterranean coastal waters using the seagrass *Posidonia oceanica*



J. Richir ^{a, b, *}, M. Salivas-Decaux ^c, C. Lafabrie ^{c, d}, C. Lopez y Royo ^c, S. Gobert ^a, G. Pergent ^c, C. Pergent-Martini ^c

d UMR 241 EIO, University of French Polynesia, BP 6570, 98 702 Faa'a, Tahiti, French Polynesia



^a Laboratory of Oceanology, MARE Centre, University of LIEGE, B6C, 4000 LIEGE, Sart Tilman, Belgium

^b Institute of Marine Sciences, University of Portsmouth, Ferry Road, Portsmouth PO4 9LY, United Kingdom

^c FRES 3041, EqEL, University of Corsica, Faculty of Sciences, BP 52, 20250 Corte, France