

# In reference to analysis of reflux as the aetiology of laryngeal dysplasia progression through a matched case-control study

We reviewed the article entitled: "Analysis of reflux as the etiology of laryngeal dysplasia progression through a matched case-control study."<sup>1</sup> The authors did not find differences in the level of pepsin, enterokinase and bilirubin in laryngeal dysplasia (LD) of patients with malignant transformation *versus* those without transformation. The involvement of reflux in the development of LD and laryngeal cancers is an important topic, and the realisation of such a study is important. However, we wish to draw attention to many points.

First, it is difficult to know whether the included patients with tissue pepsin really suffered from reflux. The detection of pepsin into the tissue means that patients had some pharyngeal reflux events the day before the surgery but cannot confirm the diagnosis. The sensitivity of pepsin detection in laryngeal tissue depends on the technique and the material (antibodies), reaching 75 to 85% depending on the type of reflux (acid vs non-acid).<sup>2</sup> Moreover, we have no detailed information about the immunostaining technique, limiting the reproducibility of the protocol. The presence of pepsin into the tissue does not ensure the reflux diagnosis. Thus, for example, it has been showed that the back flow of gastric content and the deposit of pepsin into the tissue are influenced by the meals preceding the sample collection, making the pepsin tissue a poorly reliable marker of reflux.<sup>3</sup> To improve the sensitivity, authors<sup>1</sup> could have performed hypopharyngeal-oesophageal pH-impedance monitoring, which is the only way to confirm the diagnosis.<sup>4</sup>

Second, the LD malignant transformation involves many factors such as tobacco history, environmental factors, genetic or immune response.<sup>5</sup> The authors did not provide information about the tobacco history (pack-year data) of groups, which is an important data to consider the risk of malignant transformation. Even many years after the tobacco cessation, it is conceivable that patients with long/more severe history of tobacco consumption may have more cell mucosa DNA impairments and a higher risk to develop cancer.



Third, the focus on pepsin as the only enzyme associated with malignant transformation limits the understanding of transformation mechanisms. More than 50% of patients had mixed or non-acid reflux,<sup>4</sup> in which the activity of pepsin is decreased regarding the alkaline pH of refluxate. To reliably investigate the involvement of reflux in the malignant transformation, authors have to consider the entire content of refluxate, including bile salts and trypsin.<sup>4</sup> Furthermore, bile salts may be involved in laryngopharyngeal malignant transformation.<sup>6</sup>

In future studies, reflux has to be diagnosed at the LD diagnosis time and physicians have to follow the reflux clinical course over the

time. More than 50% of reflux patients had chronic course,<sup>4</sup> which leads to a potential higher risk to develop cell DNA damage and lesions. Thus, cross-sectional study design is probably not adequate to study a disease association involving chronic and repeated exposure.

## CONFLICT OF INTEREST

None to declare.

Jerome R. Lechien<sup>1,2,3</sup>   
 Petros D. Karkos<sup>4</sup>   
 Sven Saussez<sup>1,2</sup>  
 Stephane Hans<sup>1,3</sup>

<sup>1</sup>Department of human Anatomy and Experimental Oncology, Mons School of Medicine, UMONS Research Institute for Health Sciences and Technology, University of Mons (UMons), Mons, Belgium

<sup>2</sup>Department of Otorhinolaryngology-Head and Neck Surgery, CHU Saint-Pierre, School of Medicine, Université Libre de Bruxelles, Belgium

<sup>3</sup>Department of Otorhinolaryngology-Head and Neck Surgery, Foch Hospital, UVSQ, Paris Saclay University, Paris, France

<sup>4</sup>Department of Otorhinolaryngology and Head and Neck Surgery, Thessaloniki Medical School, Thessaloniki, Greece

## Correspondence

Jerome R. Lechien, Laboratory of Anatomy and Cell Biology, Faculty of Medicine, University of Mons (UMONS), Avenue du Champ de mars, 6, B7000 Mons, Belgium.  
 Email: Jerome.Lechien@umons.ac.be

## ORCID

Jerome R. Lechien  <https://orcid.org/0000-0002-0845-0845>  
 Petros D. Karkos  <https://orcid.org/0000-0003-2488-5539>

## REFERENCES

- Gu W, Niu YY, Wang WZ, et al. Analysis of reflux as the aetiology of laryngeal dysplasia progression through a matched case-control study. *Clin Otolaryngol*. 2021;46(1):175-180. <https://doi.org/10.1111/coa.13638>
- Jiang A, Liang M, Su Z, et al. Immunohistochemical detection of pepsin in laryngeal mucosa for diagnosing laryngopharyngeal reflux. *Laryngoscope*. 2011;121(7):1426-1430. <https://doi.org/10.1002/lary.21809>

3. Lechien JR, Bobin F, Muls V, et al. Saliva pepsin concentration of laryngopharyngeal reflux patients is influenced by meals consumed before the samples. *Laryngoscope*. 2020. <https://doi.org/10.1002/lary.28756>. [Epub ahead of print].
4. Lechien JR, Bock JM, Carroll TL, Akst LM. Is empirical treatment a reasonable strategy for laryngopharyngeal reflux? A contemporary review. *Clin Otolaryngol*. 2020;45(4):450-458. <https://doi.org/10.1111/coa.13518>
5. Trivedi S, Rosen CA, Ferris RL. Current understanding of the tumor microenvironment of laryngeal dysplasia and progression to invasive cancer. *Curr Opin Otolaryngol Head Neck Surg*. 2016;24(2):121-127. <https://doi.org/10.1097/MOO.0000000000000245>
6. Shellman Z, Aldhahrani A, Verdon B, et al. Bile acids: a potential role in the pathogenesis of pharyngeal malignancy. *Clin Otolaryngol*. 2017;42(5):969-973. <https://doi.org/10.1111/coa.12822>

**How to cite this article:** Lechien JR, Karkos PD, Saussez S, Hans S. In reference to analysis of reflux as the aetiology of laryngeal dysplasia progression through a matched case-control study. *Clin Otolaryngol*. 2021;46:687–688. <https://doi.org/10.1111/coa.13711>