



EVENT ABSTRACT

A 3D neurons model to evaluate the neurotoxicity of harmane, a β carboline alkaloid incriminated in Essential Tremor

Rania Aro^{1*}, Alpha Diallo^{2, 3}, Marvyn pichueque^{2, 3}, Mathilde Wauters⁴, Amandine Nachtergaele¹, Laurence Ris⁴, Pierre Duez¹ and Mario Manto^{2, 3}

¹ University of Mons, Belgium

² Free University of Brussels, Belgium

³ Service de Neurologie, Centre Hospitalier Universitaire de Charleroi, Belgium

⁴ Research Institute for Biosciences, University of Mons, Belgium

Introduction:

β -Carbolines alkaloids (β CAs) are indole alkaloids, such as harmane, that naturally occur in plants and food and are also endogenously produced in mammals and humans. Some β CAs are potent tremor inductors and exhibit neurotoxicological activities. They have been incriminated in the pathogenesis of Essential tremor (ET) as blood Harmane levels have been found to be abnormally increased in ET patients. Harmane has also been detected in post-mortem brain samples. Essential tremor (ET) is one of the most prevalent neurological diseases but, despite a high prevalence, its physiopathology remains poorly understood. There is converging evidence that both genetic and environmental factors play a role in ET pathogenesis and so the identification of involved environmental factors is a critical step towards risk reduction through the implementation of prevention strategies or interventional studies. 2D cell cultures do not represent the real cell environment where cells spatially and chemically interact; their lack in predictivity increases the cost and failure rate of clinical trials, especially in neurosciences. Recently, 3D cell cultures have received much attention, as these are closer to tissues models, although their main challenges reside in end-point measurements.

Methods:

We propose the use of a 3D neural spheroid cultures ("mini-brains") in vitro model to assess the neurotoxic effects of harmane. We have generated a model of 3D spheroids from embryonic mouse cortical neurons using micro-molded agarose wells. After exposure to harmane at increasing concentrations of 50, 100, 250 μ M, a resazurin assay was performed to measure cell viability and a highly sensitive fluorometric method, based on the oxidation of dichlorodihydrofluorescein DCFH, was used to measure eventually induced reactive oxygen species (ROS).

Results:

Hydrogel microwells facilitated the assembly of spheroids containing neurons and glia into a complex 3D structures and prevented the agglomeration of spheroids. Exposure to harmane induced a cytotoxicity in 3D neural spheroids that was correlated with harmane concentrations, with a viability down to 24 % at 250 μ M. On the other hand, lower levels of oxidative stress were detected as the harmane concentration increased, although the effect was limited in time.

Conclusions:

We demonstrated our ability to produce a 3D culture using simple materials and routine laboratory equipment. Despite high challenges related to 3D cultures and data acquisition, this 3D neural spheroid model mimics a neuronal microenvironment, allowing a fine study of neurodegenerative disorder pathologies and the effect of chemicals on the brain. The viability and oxidative stress data support so far a dual role of harmane, as both neurotoxic and neuroprotective agent. Further studies are required to elucidate its biological activities and their eventual relationship with the ET disorder.

Acknowledgements

This research is in part funded through a FMRH grant and a Kangaroo grant from the UMONS Health Institute .

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- [1].Lavita SI, Aro R, Kiss B, Manto M, Duez P. The Role of beta-Carboline Alkaloids in the Pathogenesis of Essential Tremor. *Cerebellum*. 2016;15(3):276-84. [2].Louis ED, Zheng W. Beta-carboline alkaloids and essential tremor: exploring the environmental determinants of one of the most prevalent neurological diseases. *ScientificWorldJournal*. 2010;10:1783-94. [3].Duval K, Grover H, Han L-H, Mou Y, Pegoraro AF, Fredberg J, et al. Modeling Physiological Events in 2D vs. 3D Cell Culture. *Physiology*. 2017;32(4):266-77. [4].Bonnier F, Keating ME, Wrobel TP, Majzner K, Baranska M, Garcia-Munoz A, et al. Cell viability assessment using the Alamar blue assay: a comparison of 2D and 3D cell culture models. *Toxicology In Vitro*. 2015;29(1):124-31. [5].Dingle Y-TL, Boutin ME, Chirila AM, Livi LL, Labriola NR, Jakubek LM, et al. Three-Dimensional Neural Spheroid Culture: An In Vitro Model for Cortical Studies. *Tissue Engineering Part C, Methods*. 2015;21(12):1274-83. [6].Pamies D, Block K, Lau P, Gribaldo L, Pardo CA, Barreras P, et al. Rotenone exerts developmental neurotoxicity in a human brain spheroid model. *Toxicology and Applied Pharmacology*. 2018;354:101-14.

Keywords: Essential Tremor, 3D spheroid culture, B-Carboline alkaloids, Harmane, Toxicity

Conference: 13th National Congress of the Belgian Society for Neuroscience . Brussels, Belgium, 24 May - 24 May, 2019.

Presentation Type: Poster presentation**Topic:** Cellular/Molecular Neuroscience

Citation: Aro R, Diallo A, Pichueque M, Wauters M, Nachtergaele A, Ris L, Duez P and Manto M(2019). A 3D neurons model to evaluate the neurotoxicity of harmane, a B carboline alkaloid incriminated in Essential Tremor. *Front. Neurosci. Conference Abstract: 13th National Congress of the Belgian Society for Neuroscience* . doi:10.3389/conf.fnins.2019.96.00076

* **Correspondence:** Ms. Rania Aro, , University of Mons, Mons, Belgium, rania.aro@student.umons.ac.be

[< Back](#)

[Back to top](#)

[Home](#)

[About Frontiers](#)

[Journals A-Z](#)

[Institutional Membership](#)

[Contact](#)

[Media Relations](#)

[News](#)

[Blog](#)

[Submit](#)

[FAQs](#)

[Terms & Conditions](#)

[Newsletters](#)

[RSS/Twitter](#)

[Team](#)

[Careers](#)