Chiral Higher Spin Gravity and Convex Geometry*

Alexey Sharapov, ¹ Evgeny Skvortsov, ² and Richard Van Dongen ²

¹Physics Faculty, Tomsk State University, Lenin ave. 36, Tomsk 634050, Russia

²Service de Physique de l'Univers, Champs et Gravitation,

Université de Mons, 20 place du Parc, 7000 Mons, Belgium †

(Dated: January 13, 2023)

Chiral Higher Spin Gravity is the minimal extension of the graviton with propagating massless higher spin fields. It admits any value of the cosmological constant, including zero. Its existence implies that Chern–Simons vector models have closed subsectors and supports the 3d bosonization duality. In this letter, we explicitly construct an A_{∞} -algebra that determines all interaction vertices of the theory. The algebra turns out to be of pre-Calabi–Yau type. The corresponding products, some of which originate from Shoikhet–Tsygan–Kontsevich formality, are given by integrals over the configuration space of convex polygons.

INTRODUCTION

The idea of Higher Spin Gravity (HiSGRA) is to construct quantum gravity models by exploring extensions of gravity with massless fields of all spins [1]. While masslessness can simulate the high energy regime, the importance of higher spin states for the quantum gravity problem is supported by string theory, whose spectrum is populated by massive higher spin fields, and by the AdS/CFT correspondence since conformal field theories in $d \geq 3$ have single-trace operators of arbitrarily high spin, e.g. (Chern–Simons) vector models and $\mathcal{N}=4$ SYM [2]. Even before constructing any HiSGRA it can be seen that the smallest multiplet is unbounded in spin with an ∞ -dimensional symmetry behind [3–5].

Due to HiSGRA being at the brink it should not be surprising that constructing such theories faces numerous problems and naive attempts come immediately in contradiction with the basic field theory requirements [6–12]. Within AdS/CFT correspondence HiSGRA should be dual to vector models [13–16] or to (free) $\mathcal{N}=4$ SYM [17–19], which is based on matching the spectrum of fields/operators. However, not every CFT has a well-defined, (i) quasi-classical, (ii) local dual with (iii) finitely-many fields. HiSGRA duals of vector models immediately violate (ii) and (iii). It is (ii) that presents a serious obstacle, eliminating the standard field theory tools in an attempt to construct the theories [20].

With all hurdles already mentioned, it comes as a surprise that there exists a class of well-defined, local HiS-GRA for any value of the cosmological constant, including zero. Its flat space origin is easier to explain. As is well-known [21, 22], there is a unique cubic amplitude/vertex for any three given helicities $\lambda_1 + \lambda_2 + \lambda_3 > 0$:

$$V_{\lambda_1,\lambda_2,\lambda_3} \sim [12]^{\lambda_1+\lambda_2-\lambda_3} [23]^{\lambda_2+\lambda_3-\lambda_1} [13]^{\lambda_1+\lambda_3-\lambda_2}, (1)$$

and its complex conjugate is expressed in terms of $\langle ij \rangle$ and is valid for $\lambda_1 + \lambda_2 + \lambda_3 < 0$. Each theory comes with a particular spectrum of helicities and with cubic

(and possibly higher) couplings $C_{\lambda_1,\lambda_2,\lambda_3}$,

$$V_3 = \sum_{\lambda_1, \lambda_2, \lambda_3} C_{\lambda_1, \lambda_2, \lambda_3} V_{\lambda_1, \lambda_2, \lambda_3}. \tag{2}$$

Chiral HiSGRA [23–25] is a unique class of theories that completes a genuine higher spin interaction [26] to a Lorentz-invariant, local theory. The spectrum has to contain all helicities (at least even) and the coupling constants are uniquely fixed to be [23–25]

$$C_{\lambda_1,\lambda_2,\lambda_3} = \frac{\kappa (l_p)^{\lambda_1 + \lambda_2 + \lambda_3 - 1}}{\Gamma(\lambda_1 + \lambda_2 + \lambda_3)},$$
 (3)

where l_p is of length dimension. No higher order interactions are needed. All tree-level amplitudes can be shown to vanish on-shell (like in self-dual Yang-Mills theory and self-dual gravity) and the theory is at least one-loop finite [27–29]. These results were obtained in the lightcone gauge and argued to admit a smooth deformation to $(A)dS_4$, which was supported by [30, 31].

Chiral HiSGRA's spectrum suggests the dual to be (Chern–Simons) vector models. However, the interactions being chiral, it has to be dual to a closed subsector thereof [25, 31, 32]. Through the ABJ triality [33] one can also argue that its supersymmetric $\mathcal{N}=6$ version is dual to a closed subsector of tensionless strings on $AdS_4 \times \mathbb{CP}^3$ [34]. Chiral HiSGRA can also be detected through celestial studies [35, 36].

The problem of covariantization and extension to $(A)dS_4$ of Chiral HiSGRA was solved in the recent papers [32, 37, 38]. However, the solution was wrapped into a rather abstract form of homological perturbation theory with considerable technical difficulties to extract explicit vertices and with many hidden gems that we uncover in this letter.

The main result of the letter is an explicit construction of classical field equations for Chiral HiSGRA. These originate from a cyclic A_{∞} -algebra of pre-Calabi–Yau type [39]. The (higher) multiplications of the A_{∞} -algebra are given by integrals over the configuration space of some convex polygons. This gives the first example of a

local covariant HiSGRA with propagating massless fields. A grain of salt is that Chiral HiSGRA appears to be non-unitary due to interactions being complex in Minkowski signature and it is close in spirit to self-dual theories. Nevertheless, the theory should be unitary in flat space [27-29] at the price of having the trivial S-matrix, while in $(A)dS_4$, similarly to self-dual theories, the fact that it is a closed subsector of a unitary theory implies that all solutions and amplitudes of Chiral theory should carry over to the holographic dual of (Chern–Simons) vector models.

CHIRAL HISGRA AND A_{∞}

Initial higher spin data

For a given set of physical degrees of freedom there is more than one way to incorporate them into Lorentz covariant (spin-)tensor fields. For a massless spin-s particle a common way is to take a rank-s symmetric tensor $\Phi_{\mu_1\cdots\mu_s}$, known as Fronsdal's field [40]. In the spinorial language its traceless component takes values in the (s, s)-representation of the Lorentz algebra [41], $\Phi_{A(s),A'(s)}$. In principle, any spin-tensor $\Phi_{A(n),A'(m)}$ with n+m=2s allows one to describe the same degrees of freedom. Most of these spin-tensor fields, while completely equivalent at the free level, resist including many important interactions, e.g. the gravitational one. For Chiral HiSGRA an optimal way to describe a massless spin-s particle is rooted in the twistor approach [42–45]. Accordingly, positive and negative helicity states are placed into two different fields: the zero-form $\Psi^{A(2s)}$ and the one-form $\Omega^{A(2s-2)}$. The free action reads [46]

$$S = \int \Psi^{A(2s)} \wedge e_{AB'} \wedge e_{A}^{B'} \wedge \nabla \Omega_{A(2s-2)}, \quad (4)$$

where $e^{AA'}$ is the vierbein one-form and ∇ is the compatible Lorentz-covariant derivative ($\nabla e^{AA'}=0$). Note that ∇ does not have to be flat/(A)dS and describes an arbitrary self-dual background, i.e., $\nabla^2 \chi^A \equiv 0$ for any χ^A . The action is invariant under the infinitesimal gauge transformations $\delta \Psi^{A(2s)}=0$ and

$$\delta\Omega^{A(2s-2)} = \nabla\xi^{A(2s-2)} + e^{A}{}_{C'}\,\eta^{A(2s-3),C'}\,, \qquad (5)$$

where the gauge parameters ξ 's and η 's are zero-forms. One can also append this action with Yang–Mills and gravitational interactions among the higher spin fields to construct higher spin extensions of self-dual Yang–Mills theory and of self-dual gravity [46, 47]. These two theories are consistent truncations of Chiral HiSGRA [47], from which the scalar field can be safely discarded for the gravitational and Yang–Mills interactions are not much restrictive. Once a genuine higher spin interaction is present a unique completion gives Chiral HiSGRA.

The fields $\Omega^{A(2s-2)}$ can be packaged into a single gauge field $\Omega(y) = \sum_{s} \Omega^{A(2s-2)} y_A \cdots y_A$ assuming its values in the Weyl algebra A_{λ} . By definition, A_{λ} is generated by the formal variables \hat{y}_A subject to the canonical commutation relations $[\hat{y}_A, \hat{y}_B] = -2\lambda \epsilon_{AB}$. We prefer to think of the algebra A_{λ} as resulting from the deformation quantization of \mathbb{R}^2 , i.e., as the space of complex polynomials in y_A endowed with the Moyal–Weyl \star -product. The numerical parameter λ will be proportional to the square root $\sqrt{|\Lambda|}$ of the cosmological constant Λ . Likewise, after extending with the scalar field, the fields $\Psi^{A(2s)}$ can be packaged into a single matter field $\Psi(y)$. It is convenient to think of $\Psi(y)$ as taking values in the dual space $A_{\lambda}^* \simeq \mathbb{C}[[y_A]],$ the space of formal power series in y's with complex coefficients. In the commutative/flat limit $\lambda = 0$, the Weyl algebra A_{λ} passes smoothly to the commutative algebra of complex polynomials $A_0 = \mathbb{C}[y_A]$.

Sigma-model

The equations of motion of Chiral HiSGRA are constructed in the (AKSZ) sigma-model form [48]:

$$d\Phi = \sum_{n=2}^{\infty} l_n(\underline{\Phi, \dots, \Phi}). \tag{6}$$

Here Φ is a collection of zero- and one-form fields, d is the exterior differential, and the l_n 's are exterior polynomials in the form fields Φ . The formal integrability of system (6), steming from $d^2 = 0$, imposes an infinite sequence of quadratic relations on the multi-linear functions l_n , which can be recognized as defining relations of a minimal L_{∞} -algebra \mathbb{L} [49]. System (6) admits a consistent truncation setting $l_n = 0$ for all n > 2. The remaining bilinear function l_2 defines then a graded Lie bracket on the target space of form fields Φ . In general, the trilinear function l_3 is defined by a Chevalley–Eilenberg cocycle of the graded Lie algebra with bracket l_2 .

In principle, any system of PDE can be cast into the form (6) at the expense of introducing auxiliary fields [50–52]. However, to describe propagating degrees of freedom the total number of components of the fields Φ must necessarily be infinite.

A remarkable fact is that the algebra \mathbb{L} underlying Chiral HiSGRA originates from a minimal A_{∞} -algebra $\hat{\mathbb{A}}$ via the standard symmetrization procedure [53]. The A_{∞} -algebra $\hat{\mathbb{A}}$ decomposes further into the tensor product $\mathbb{A} \otimes B$ of a smaller A_{∞} -algebra \mathbb{A} and a unital associative algebra B. For Chiral HiSGRA B is chosen to be $A_{\lambda=1} \otimes \operatorname{Mat}_N$. The factor A_1 is needed to have the right set of auxiliary fields for (6) to reproduce free equations of motion resulting from (4) [54] and Mat_N accommodates possible Yang–Mills gaugings. Since \mathbb{A} and B are completely independent of each other, we may keep the latter as an arbitrary parameter of $\hat{\mathbb{A}}$. It is also

striking that the A_{∞} -algebra $\mathbb A$ is of a very special type known as a pre-Calabi–Yau algebra of degree two [39]. This is defined as an A_{∞} -algebra built on $A[-1] \oplus A^*$, where $A = \bigoplus A_n$ is a graded associative algebra, A^* is its dual bimodule, and $A[-1]_n = A_{n-1}$. The A_{∞} -products m_n in $A[-1] \oplus A^*$ together with the natural pairing $\langle a|c \rangle = -\langle c|a \rangle$ for $a \in A$ and $c \in A^*$ give rise to the multi-linear forms $\langle m_n(\alpha_0, \ldots, \alpha_{n-1}) | \alpha_n \rangle$ on $A[-1] \oplus A^*$ that are required to have cyclic symmetry. In our case, $\mathbb A$ is concentrated in degrees 0 and 1, i.e., $\mathbb A = \mathbb A_0 \oplus \mathbb A_1$, where $\mathbb A_0 = A_\lambda^*$ and $\mathbb A_1 = A_\lambda[-1]$. All the products m_n in $\mathbb A$ are of degree -1. The canonical projection of the field Φ onto the A_{∞} -subalgebra $\mathbb A \subset \hat{\mathbb A}$ reproduces the pair of form fields Ω and Ψ that we started with.

Configuration space and Convex Geometry

The products m_n in the A_{∞} -algebra \mathbb{A} are reminiscent of (Shoikhet–Tsygan)–Kontsevich formality [55, 56]: they are poly-differential operators defined through integrals over configuration spaces of points inside and on the boundary of disk. Let us, however, stress that an extension of the formality that would generate the vertices of Chiral theory has not yet been identified. Nevertheless, one would expect that since the Poisson structure behind the Weyl algebra A_{λ} is ϵ_{AB} , i.e., constant and symplectic, the configuration space lacks the bulk part and reduces to points on the boundary. Therefore, the configuration space defined below should be identified with the boundary part of a yet to be found extension of the formality.

Our configuration space V_n admits two equivalent definitions: (a) the space of convex polygons that can be inscribed into a unit square and have one edge along the diagonal (region B in Fig.1) or (b) the space of all concave polygons with three convex vertices on the two adjacent sides of the unit square (region A), which we call swallowtails. Clearly, V_n is a compact region in \mathbb{R}^{2n} .

Both the definitions suffice for practical applications, but definition (b) can be improved. In a few words, one can consider the space of concave polygons with exactly three consecutive convex vertices on the oriented Euclidian plane \mathbb{E}^2 modulo the affine transformations of $GL^+(2,\mathbb{R})\ltimes\mathbb{R}^2$. Any such polygon corresponds to a string of vectors $Q=(\vec{b_1},\vec{b_2},\vec{a_1},\ldots,\vec{a_n},\vec{a_0})$ with $\vec{b_{1,2}}$ connecting the three convex vertices. Their affine coordinates form a $2\times(n+3)$ -matrix Q with vanishing sum of columns (the polygon is closed). By $GL^+(2,\mathbb{R})$ -transformations one can bring Q into the canonical form

$$Q = \begin{pmatrix} -1 & 0 & u_1 & u_2 & \dots & u_n & 1 - \sum u_i \\ 0 & -1 & v_1 & v_2 & \dots & v_n & 1 - \sum v_i \end{pmatrix}$$
 (7)

with one of the convex vertices at the origin. Let q_{ij} denote the determinant of a 2×2 minor of Q built from columns i and j. Then the sum $|Q| = \sum_{i < j} q_{ij}$ is equal




FIG. 1. Configuration space: a convex polygon B and a swallowtail A. The swallowtail is built on $-\vec{e}_1, -\vec{e}_2, \vec{a}_1, \dots, \vec{a}_n, \vec{a}_0$.

to twice the area of the swallow tail A. |Q| depends on Q modulo cyclic permutations of the columns.

The configuration space V_n can also be understood as a subset in the oriented Grassmannian $\widetilde{Gr}(2, n)$, with q_{ij} being the corresponding Plücker's coordinates [57]. It is not a positive one [58, 59], but the signs of q_{ij} are fixed.

A_{∞} -maps/Vertices

Now we are ready to construct the (higher) products $m_n(\alpha_1, \ldots, \alpha_n)$ for $\alpha_i \in \mathbb{A}$. As mentioned above they are given by poly-differential operators

$$m_n(p_0, p_1, \dots, p_n) \alpha_1(y_1) \cdots \alpha_n(y_n) \Big|_{y_i=0}$$
, (8)

where $p_0 \equiv y$ and $p_i \equiv \partial_{y_i}$. The products respect Sp(2)-symmetry and are functions of the Sp(2)-invariant scalar products $p_{ij} \equiv -\epsilon_{AB}p_i^Ap_j^B$ such that $\exp[p_{0i}]f(y_i) = f(y+y_i)$ is the shift operator. In particular, the pairing between elements $a \in A_{\lambda}$ and $c \in A_{\lambda}^*$ is defined as

$$\langle a|c\rangle = -\langle c|a\rangle = e^{p_{12}} a(y_1)c(y_2)|_{y_i=0}.$$
 (9)

Thanks to the cyclicity of the A_{∞} -structure on \mathbb{A} many products are related to each other. At order n one has $1+\left\lceil\frac{n-1}{2}\right\rceil$ independent products. By dimensional considerations, each product m_n may have either one or two arguments of \mathbb{A}_1 . By cyclicity, the products with one argument are expressed through the products with two arguments. Therefore, we describe only those products m_{n+2} that have two arguments a, b of \mathbb{A}_1 and the other n arguments c_1, \ldots, c_n , of \mathbb{A}_0 . Such products are represented by sums over all planar rooted trees with exactly two branches, see Fig. 2. Each branch ends by one of the two arguments a, b. The other leaves are decorated by the arguments c_i . It is also convenient to decorate the root of the tree with an additional argument c_0 , which corresponds to the scalar $\langle m_{n+2}(c_1, \ldots, a, \ldots, b, \ldots, c_n) | c_0 \rangle$.




FIG. 2. A generic tree T on the left and the only tree T_0 contributing to (11) on the right. The counterclockwise arrow orders the arguments and i of c_i on the left panel corresponds to its position on T_0 as obtained via flips and shifts.

Let us start with the simplest tree T_0 , see the right panel in Fig. 2. With each c_i and c_0 we associate two vectors: $\vec{a}_i = (u_i, v_i)$, i = 1, ..., n, and $\vec{a}_0 = (1 - \sum_i u_i, 1 - \sum_i v_i)$, so that $\sum_i \vec{a}_i = 0$; $\vec{r}_i = (p_{a,i}, p_{b,i})$, i = 1, ..., n, and $\vec{r}_0 = (p_{0,a}, p_{0,b})$. This notation corresponds to the symbol of an operator that acts on

$$a(y_a)b(y_b)c_1(y_1)\cdots c_n(y_n)|_{y_{\bullet}=0}$$
. (10)

We also need an auxiliary matrix $P = (\vec{0}, \vec{0}, \vec{r}_1, \dots, \vec{r}_n, \vec{r}_0)$ and recall that $Q = (-\vec{e}_1, -\vec{e}_2, \vec{a}_1, \dots, \vec{a}_n, \vec{a}_0)$.

The tree T_0 describes a single contribution to the product $m_{n+2}(a, b, c_1, \ldots, c_n)$ with $a, b \in \mathbb{A}_1$ and $c_i \in \mathbb{A}_0$. The corresponding symbol reads

$$m_{n+2} = (p_{a,b})^n \int_{\mathbb{V}_n} \exp\left(\text{tr}[PQ^t] + \lambda |Q| p_{a,b}\right),$$
 (11)

see also (15). It generates many other structure maps via cyclicity. All other trees that contribute to m_{n+2} give similar expressions. We just need to adjust Q and P in accordance with the topology of a given tree. Every tree T can be obtained from T_0 by two operations: (1) flipping some c_i from right to left on the right branch; (2) shifting the string of c_1, \ldots, c_n, c_0 by one unit along the cord connecting b to a. After these operations c_i still remember their vectors \vec{a}_i and \vec{r}_i from T_0 , e.g. c_i of T in Fig. 2 has \vec{a}_i and $\vec{r_i}$ associated with it. They also remember their arguments, e.g. $c_i \equiv c_i(y_i)$ as it was for T_0 . The arguments of the symbol that corresponds to T should be read off from left to right. Therefore, the labeling of $c_i \equiv c_i(y_i)$, which is inherited from T_0 , does not correspond to the natural labeling $c_1(y_1), \ldots, a(y_a), \ldots, b(y_b), \ldots, c_n(y_n)$. This will be compensated by a permutation σ_T that reshuffles the labels on the c_i 's and p_{ij} 's. It also permutes the arguments y_i of c_i . The symbol associated with T consists of the sign, prefactor, principal term, and cosmological term (the last two are in the exponent). They are computed as follows.

Sign. For a given tree T the sign s_T is equal to $(-1)^m$, where m is the total number of c's in between a and b. Note that this number is cyclic invariant, which is necessary for the cyclicity to work.

Prefactor. The prefactor is given by $(p_{a,b})^n$ where n is the number of c's; it acts on a and b.

Principal term. To construct P_T for a given tree T one can just use the cyclicity applied to T_0 . Explicitly, $P_T = (\vec{0}, \vec{0}, \vec{r_1}, \dots, -r_j, \dots, \vec{r_n}, -\vec{r_0})$, where j is the c_j at the root of T. The principal term is $\operatorname{tr}[P_T Q^t]$.

Cosmological term. We associate the first two vectors $-e_1$, $-e_2$ of Q (7) with a and b. Recall that vectors \vec{a}_i are associated with c_i , including c_0 . In order to construct Q_T for a given tree T we fill in the columns of Q_T starting from a and then following the tree counterclockwise. The coefficient of λ is $|Q_T| p_{a,b}$; it acts on a and b.

Now we combine all the ingredients together and apply the permutation σ_T to bring the labeling inherited from T_0 into the natural one. Thus, each tree T makes the following contribution to m_{n+2} :

$$s_T \sigma_T(p_{a,b})^n \int_{\mathbb{V}_n} \exp\left(\operatorname{tr}[P_T Q^t] + \lambda |Q_T| p_{a,b}\right). \tag{12}$$

One needs to sum over all trees T to construct all m_{n+2} . We claim [60] that the products constructed in such a way do satisfy the defining relations of an A_{∞} -algebra. Combining these products with the associative product in B, we can write the r.h.s. of equation (6) as

$$l_n(\Phi, \dots, \Phi) = m_n(\Phi, \dots, \Phi) \tag{13}$$

for the form field Φ with values in $\mathbb{A}=\mathbb{A}\otimes B$. When restricted to the Φ -diagonal, symmetrization is automatically performed, turning the A_{∞} -algebra products into the multi-brackets l_n of the L_{∞} -algebra \mathbb{L} .

An important property of the poly-differential operators m_n is that the corresponding symbols do not involve p_{ij} 's that connect c_i with c_j . This translates into the locality of the vertices in the field theory language. Another important property is that the flat space vertices smoothly deform to $(A)dS_4$. In other words, the flat space limit is nonsingular.

Low order products

By way of illustrations let us present some low order products of \mathbb{A} . For m_2 the configuration space \mathbb{V}_0 is zero-dimensional and the corresponding swallowtail occupies half of the square (area = 1/2). The associated symbol

$$m_2(a,b) = \exp[p_{0,a} + p_{0,b} + \lambda p_{a,b}]$$
 (14)

is just the Moyal–Weyl *-product with parameter λ . Cyclicity implies $\langle m_2(a,b)|c\rangle = \langle a|m_2(b,c)\rangle = -\langle b|m_2(c,a)\rangle$, which just gives the bimodule structure on A^*_{λ} , as designed. These vertices along reproduce most of the cubic amplitudes (1), (3) of Chiral HiSGRA [61].

For m_3 the space V_1 is two-dimensional: one can place a point (u, v) anywhere below the diagonal, the volume




FIG. 3. Configuration spaces V_1 and V_2 .

of V_1 is 1/2 and the area of A is $\frac{1}{2}(1+u-v)$, see Fig. 3. The duality determines 6 maps via just 2:

$$\langle m_3(a, b, c_1) | c_2 \rangle = \langle m_3(c_2, a, b) | c_1 \rangle ,$$

$$\langle m_3(a, b, c_1) | c_2 \rangle = \langle a | m_3(b, c_1, c_2) \rangle ,$$

$$\langle m_3(c_1, a, b) | c_2 \rangle = \langle m_3(c_2, c_1, a) | b \rangle ,$$

$$\langle m_3(a, c_1, b) | c_2 \rangle = \langle a | m_3(c_1, b, c_2) \rangle .$$

The two basis products can be chosen as $m_3(a, b, c)$ and $m_3(a, c, b)$, see (16).

For m_4 the space \mathbb{V}_2 is four-dimensional and its volume is equal to 1/24. There are two independent products: one resulting from (15) for n=2 and another one given by the sum (17) over three trees.

CONCLUSIONS AND DISCUSSION

The main results of this letter are (i) an explicit construction of all covariant interactions vertices of Chiral HiSGRA both in flat and $(A)dS_4$ spaces; (ii) a remarkable relation between the vertices and convex polygons; (iii) a rich class of 2-pre-Calabi-Yau algebras \hat{A} that are parameterized by an associative algebra with trace. This gives the first example of a well-defined, local, manifestly covariant HiSGRA with propagating massless fields [62]. The results open up many obvious directions: (a) calculation of holographic correlation functions; (b) constructing exact solutions; (c) looking for an action [63] that would covariantize the light-cone results and extend them to $(A)dS_4$. Eventually, one expects Chiral HiSGRA to be integrable and UV finite, which is still to be proved.

In the regard to item (iii) a lot needs to be understood. It is clear that the field theory underlying Chiral HiSGRA is low dimensional since the functional dimension of A_{λ} is two and the extra associative factor B in $\hat{\mathbb{A}} = \mathbb{A} \otimes B$ is a passive spectator. This should be related to an important result of [47] that in the light-cone gauge the equations

of motion can be cast into the form of the principal chiral model. Among other ideas, \mathbb{A} can be used to construct field theories in 2d and 3d. One can also construct a plenty of theories as a double-copy, just tensoring \mathbb{A} with any A_{∞} -algebra, e.g. $\mathbb{A} \otimes \mathbb{A}$.

Chiral HiSGRA is a local field theory in AdS_4 and it can be treated by the standard AdS_4/CFT_3 -tools to give correlation functions of higher spin currents on the CFT side. They should cover a closed subsector of (Chern–Simons) vector models, which is yet to be identified. Nevertheless, the very existence of such a closed subsector supports [31, 64] the 3d bosonization duality [16, 65–69].

There is also a distant relation to tensionless string theory on $AdS_4 \times \mathbb{CP}^3$, which is via the ABJ triality [33]. One needs $\mathcal{N}=6$ supersymmetric Chiral HiSGRA, which is easily achieved via the right B-factor of Clifford algebra [70]. Again, the very existence of Chiral HiSGRA implies that there should also exist a closed subsector of this tensionless string theory.

It is worth stressing that the 'formality' underlying the construction of Chiral HiSGRA is not yet known. The cubic product m_3 is equivalent [71, 72] to one that follows from the Shoikhet–Tsygan–Kontsevich formality [55, 56]. Higher structure maps are related to the deformation quantization of the Poisson orbifold $\mathbb{R}^2/\mathbb{Z}_2$ [72]. However, these relations can be seen at the formal A_{∞} -level and do not give the specific vertices of the present paper. There should exist a topological field theory behind our construction [73], which the appearance of a pre-Calabi–Yau algebra also suggests [39].

The products of the A_{∞} -algebra $\mathbb A$ underlying Chiral HiSGRA are fine-tuned to represent a local field theory. Indeed, a generic A_{∞} -automorphism would lead to a non-local field redefinition for the vertices of sigma-model (6), thereby violating the equivalence theorem. This leads us to conclude that A_{∞}/L_{∞} -algebras representing actual field theories must have preferred bases where the vertices are maximally local. It would be interesting to reformulate the property of 'maximal locality' in a pure algebraic form.

Lastly, it is worth mentioning other interesting examples of HiSGRA: 3d topological [74–80]; 4d (and all even dimensions) conformal HiSGRA [81–83]; twistor constructions [63, 84–86]; IKKT-based [86–88]; holographic reconstruction [89, 90]. The last two examples relax the locality assumption in a controllable way and are not, strictly speaking, field theories. It seems even necessary to go beyond the field theory approach to construct HiSGRA's with massless fields that extend and complete Chiral HiSGRA.

$$m_{n+2}(a,b,c_1,\ldots,c_n) = (p_{a,b})^n \int_{\mathbb{V}_n} \exp\left[(1 - \sum_i u_i) p_{0,a} + (1 - \sum_i v_i) p_{0,b} + \sum_i u_i p_{a,i} + \sum_i v_i p_{b,i} + \lambda \left(1 + \sum_i (u_i - v_i) + \sum_{i,j} u_i v_j \operatorname{sign}(j-i) \right) p_{a,b} \right].$$
(15)

$$m_{3}(a,b,c) = +p_{a,b} \int_{\mathbb{V}_{1}} \exp[(1-u) p_{0,a} + (1-v) p_{0,b} + u p_{a,1} + v p_{b,1} + \lambda (1+u-v) p_{a,b}],$$

$$m_{3}(a,c,b) = -p_{a,b} \int_{\mathbb{V}_{1}} \exp[u p_{0,a} + v p_{0,b} + (1-u) p_{a,1} - \lambda p_{a,b} (1-u-v) - (1-v) p_{b,1}]$$

$$-p_{a,b} \int_{\mathbb{V}_{1}} \exp[v p_{0,a} + u p_{0,b} + (1-v) p_{a,1} - \lambda p_{a,b} (1-u-v) - (1-u) p_{b,1}].$$
(16)

$$m_{4}(a, c_{1}, b, c_{2}) = -p_{a,b}^{2} \int_{\mathbb{V}_{2}} \exp[(1 - u_{1} - u_{2}) p_{0,a} + (1 - v_{1} - v_{2}) p_{0,b} + u_{2} p_{a,1} + \lambda A_{1} p_{a,b} + u_{1} p_{a,2} - v_{2} p_{1,b} + v_{1} p_{b,2}] +$$

$$- p_{a,b}^{2} \int_{\mathbb{V}_{2}} \exp[(1 - u_{1} - u_{2}) p_{0,a} + (1 - v_{1} - v_{2}) p_{0,b} + u_{1} p_{a,1} + \lambda A_{2} p_{a,b} + u_{2} p_{12} - v_{1} p_{1,b} + v_{2} p_{b,2}] +$$

$$- p_{a,b}^{2} \int_{\mathbb{V}_{2}} \exp[u_{2} p_{0,a} + v_{2} p_{0,b} + (1 - u_{1} - u_{2}) p_{a,1} - \lambda A_{3} p_{a,b} + u_{1} p_{a,2} - (1 - v_{1} - v_{2}) p_{1,b} + v_{1} p_{b,2}],$$

$$A_{1} = 1 - u_{1} v_{2} + u_{2} v_{1} + u_{1} - u_{2} - v_{1} - v_{2}, \quad A_{2} = 2(1 - v_{1} - v_{2}) - A_{1}, \quad A_{3} = A_{1} - 2u_{1}.$$

$$(17)$$

- The work of E. S. and R. van D. was partially supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 101002551) and by the Fonds de la Recherche Scientifique — FNRS under Grant No. F.4544.21. A. Sh. gratefully acknowledges the financial support of the Foundation for the Advancement of Theoretical Physics and Mathematics "BASIS". E.S. expresses his gratitude to the Erwin Schrödinger Institute in Vienna for the hospitality during the program "Higher Structures and Field Theory" while this work was in progress. The results on the configuration space and Grassmannians were obtained under exclusive support of the Ministry of Science and Higher Education of the Russian Federation (project No. FSWM-2020-0033). Also at Lebedev Institute of Physics, Leninsky ave. 53, 119991 Moscow, Russia
- X. Bekaert, N. Boulanger, A. Campoleoni, M. Chiodaroli, D. Francia, M. Grigoriev, E. Sezgin, and E. Skvortsov, Snowmass White Paper: Higher Spin Gravity and Higher Spin symmetry, (2022), arXiv:2205.01567 [hep-th].
- [2] There can be regions in the coupling space where, as in the SYM example, the higher-spin states decouple and the stress-tensor multiplet remains, but only in the first approximation.
- [3] M. Flato and C. Fronsdal, One Massless Particle Equals Two Dirac Singletons: Elementary Particles in a Curved Space. 6., Lett.Math.Phys. 2, 421 (1978).

- [4] E. S. Fradkin and M. A. Vasiliev, Candidate to the role of higher spin symmetry, Ann. Phys. 177, 63 (1987).
- [5] N. Boulanger, D. Ponomarev, E. D. Skvortsov, and M. Taronna, On the uniqueness of higher-spin symmetries in AdS and CFT, Int. J. Mod. Phys. A28, 1350162 (2013), arXiv:1305.5180 [hep-th].
- [6] S. Weinberg, Photons and Gravitons in S Matrix Theory: Derivation of Charge Conservation and Equality of Gravitational and Inertial Mass, Phys. Rev. 135, B1049 (1964).
- [7] S. R. Coleman and J. Mandula, All Possible Symmetries of the S Matrix, Phys. Rev. 159, 1251 (1967).
- [8] A. Fotopoulos and M. Tsulaia, On the Tensionless Limit of String theory, Off - Shell Higher Spin Interaction Vertices and BCFW Recursion Relations, JHEP 11, 086, arXiv:1009.0727 [hep-th].
- [9] X. Bekaert, J. Erdmenger, D. Ponomarev, and C. Sleight, Quartic AdS Interactions in Higher-Spin Gravity from Conformal Field Theory, JHEP 11, 149, arXiv:1508.04292 [hep-th].
- [10] J. Maldacena, D. Simmons-Duffin, and A. Zhiboedov, Looking for a bulk point, JHEP 01, 013, arXiv:1509.03612 [hep-th].
- [11] C. Sleight and M. Taronna, Higher-Spin Gauge Theories and Bulk Locality, Phys. Rev. Lett. 121, 171604 (2018), arXiv:1704.07859 [hep-th].
- [12] D. Ponomarev, A Note on (Non)-Locality in Holographic Higher Spin Theories, Universe 4, 2 (2018), arXiv:1710.00403 [hep-th].
- [13] I. R. Klebanov and A. M. Polyakov, AdS dual of the critical O(N) vector model, Phys. Lett. **B550**, 213 (2002),

- arXiv:hep-th/0210114.
- [14] E. Sezgin and P. Sundell, Holography in 4D (super) higher spin theories and a test via cubic scalar couplings, JHEP 0507, 044, arXiv:hep-th/0305040 [hep-th].
- [15] R. G. Leigh and A. C. Petkou, Holography of the N=1 higher spin theory on AdS(4), JHEP 0306, 011, arXiv:hep-th/0304217 [hep-th].
- [16] S. Giombi, S. Minwalla, S. Prakash, S. P. Trivedi, S. R. Wadia, and X. Yin, Chern-Simons Theory with Vector Fermion Matter, Eur. Phys. J. C72, 2112 (2012), arXiv:1110.4386 [hep-th].
- [17] B. Sundborg, Stringy gravity, interacting tensionless strings and massless higher spins, Nucl. Phys. Proc. Suppl. 102, 113 (2001), arXiv:hep-th/0103247.
- [18] N. Beisert, M. Bianchi, J. F. Morales, and H. Samtleben, Higher spin symmetry and N=4 SYM, JHEP 07, 058, arXiv:hep-th/0405057 [hep-th].
- [19] M. R. Gaberdiel and R. Gopakumar, String Dual to Free N=4 Supersymmetric Yang-Mills Theory, Phys. Rev. Lett. 127, 131601 (2021), arXiv:2104.08263 [hep-th].
- [20] The results of [8–12] imply that this class of theories cannot be constructed by any Noether procedure. Nevertheless, one can 'reconstruct' them from correlation functions [9, 89, 90], which, however, has its own puzzles and is not applicable to Chern–Simons vector models before they are solved.
- [21] A. K. H. Bengtsson, I. Bengtsson, and N. Linden, Interacting Higher Spin Gauge Fields on the Light Front, Class. Quant. Grav. 4, 1333 (1987).
- [22] P. Benincasa and E. Conde, Exploring the S-Matrix of Massless Particles, Phys. Rev. D 86, 025007 (2012), arXiv:1108.3078 [hep-th].
- [23] R. R. Metsaev, Poincare invariant dynamics of massless higher spins: Fourth order analysis on mass shell, Mod. Phys. Lett. A6, 359 (1991).
- [24] R. R. Metsaev, S matrix approach to massless higher spins theory. 2: The Case of internal symmetry, Mod. Phys. Lett. A6, 2411 (1991).
- [25] D. Ponomarev and E. D. Skvortsov, Light-Front Higher-Spin Theories in Flat Space, J. Phys. A50, 095401 (2017), arXiv:1609.04655 [hep-th].
- [26] One- and two-derivative Yang-Mills and gravitational vertices are not enough. One needs higher-derivative non-abelian interaction with a higher spin field.
- [27] E. D. Skvortsov, T. Tran, and M. Tsulaia, Quantum Chiral Higher Spin Gravity, Phys. Rev. Lett. 121, 031601 (2018), arXiv:1805.00048 [hep-th].
- [28] E. Skvortsov, T. Tran, and M. Tsulaia, More on Quantum Chiral Higher Spin Gravity, Phys. Rev. D101, 106001 (2020), arXiv:2002.08487 [hep-th].
- [29] E. Skvortsov and T. Tran, One-loop Finiteness of Chiral Higher Spin Gravity, (2020), arXiv:2004.10797 [hep-th].
- [30] R. R. Metsaev, Light-cone gauge cubic interaction vertices for massless fields in AdS(4), Nucl. Phys. B936, 320 (2018), arXiv:1807.07542 [hep-th].
- [31] E. Skvortsov, Light-Front Bootstrap for Chern-Simons Matter Theories, JHEP 06, 058, arXiv:1811.12333 [hep-th].
- [32] A. Sharapov and E. Skvortsov, Chiral Higher Spin Gravity in (A)dS $_4$ and secrets of Chern–Simons Matter Theories, (2022), arXiv:2205.15293 [hep-th].
- [33] C.-M. Chang, S. Minwalla, T. Sharma, and X. Yin, ABJ Triality: from Higher Spin Fields to Strings, J. Phys. A46, 214009 (2013), arXiv:1207.4485 [hep-th].

- [34] E.S. is grateful to Andre Coimbra for asking a question that leads to this idea.
- [35] L. Ren, M. Spradlin, A. Yelleshpur Srikant, and A. Volovich, On effective field theories with celestial duals, JHEP 08, 251, arXiv:2206.08322 [hep-th].
- [36] R. Monteiro, Celestial chiral algebras, colour-kinematics duality and integrability, (2022), arXiv:2208.11179 [hep-th].
- [37] E. Skvortsov and R. Van Dongen, Minimal models of field theories: Chiral higher spin gravity, Phys. Rev. D 106, 045006 (2022), arXiv:2204.10285 [hep-th].
- [38] A. Sharapov, E. Skvortsov, A. Sukhanov, and R. Van Dongen, Minimal model of Chiral Higher Spin Gravity, (2022), arXiv:2205.07794 [hep-th].
- [39] M. Kontsevich, A. Takeda, and Y. Vlassopoulos, Pre-Calabi-Yau algebras and topological quantum field theories, arXiv preprint arXiv:2112.14667 (2021).
- [40] C. Fronsdal, Massless fields with integer spin, Phys. Rev. D18, 3624 (1978).
- [41] Hereinafter $A, B, \ldots = 1, 2, A', B', \ldots = 1, 2$ are the indices of two-fundamental representations of the Lorentz algebra, e.g. of $sl(2, \mathbb{C})$ for the Minkowski signature. We also abbreviate a group of totally symmetric (or to be symmetrized) indices $A_1 \ldots A_k$ as A(k). All spinor indices are raised and lowered with the help of the antisymmetric ϵ -symbols ϵ_{AB} and $\epsilon_{A'B'}$ with $\epsilon_{12} = 1$.
- [42] R. Penrose, Zero rest mass fields including gravitation: Asymptotic behavior, Proc. Roy. Soc. Lond. A284, 159 (1965).
- [43] L. P. Hughston, R. S. Ward, M. G. Eastwood, M. L. Ginsberg, A. P. Hodges, S. A. Huggett, T. R. Hurd, R. O. Jozsa, R. Penrose, A. Popovich, et al., eds., Advances in twistor theory (1979).
- [44] M. G. Eastwood, R. Penrose, and R. O. Wells, Cohomology and Massless Fields, Commun. Math. Phys. 78, 305 (1981).
- [45] N. M. J. Woodhouse, Real methods in twistor theory, Class. Quant. Grav. 2, 257 (1985).
- [46] K. Krasnov, E. Skvortsov, and T. Tran, Actions for Self-dual Higher Spin Gravities, (2021), arXiv:2105.12782 [hep-th].
- [47] D. Ponomarev, Chiral Higher Spin Theories and Self-Duality, JHEP 12, 141, arXiv:1710.00270 [hep-th].
- [48] It was first introduced by Sullivan [91] as a Free Differential Algebra and later leaked into supergravity [92, 93] and higher spins [94]. In modern terms this is AKSZ equations of motion [95], see [96] for the relation to the HiSGRA problem and [52] for a broader context.
- [49] T. Lada and J. Stasheff, Introduction to SH Lie algebras for physicists, Int. J. Theor. Phys. 32, 1087 (1993), arXiv:hep-th/9209099 [hep-th].
- [50] G. Barnich and M. Grigoriev, A Poincare lemma for sigma models of AKSZ type, J. Geom. Phys. 61, 663 (2011), arXiv:0905.0547 [math-ph].
- [51] G. Barnich and M. Grigoriev, First order parent formulation for generic gauge field theories, JHEP 1101, 122, arXiv:1009.0190 [hep-th].
- [52] M. Grigoriev and A. Kotov, Gauge PDE and AKSZtype Sigma Models, Fortsch. Phys. 67, 1910007 (2019), arXiv:1903.02820 [hep-th].
- [53] This is just an A_∞-extension of the familiar statement that the commutator in any associative algebra defines a Lie bracket.
- [54] This set of auxiliary fields depends on the free spectrum

- and, for that reason, is exactly the same as in [94, 97]. Some further simple projections/reality conditions may be needed, e.g. to reduce Mat_N to U(N). An additional factor of Clifford algebra will lead to supersymmetric extensions.
- [55] M. Kontsevich, Deformation quantization of Poisson manifolds. 1., Lett. Math. Phys. 66, 157 (2003), arXiv:qalg/9709040 [q-alg].
- [56] B. Shoikhet, A proof of the Tsygan formality conjecture for chains, Advances in Mathematics 179, 7 (2003).
- [57] A. Sharapov, E. Skvortsov, A. Sukhanov, and R. Van Dongen, More on Chiral Higher Spin Gravity and Convex Geometry, to appear (2022).
- [58] Nima Arkani-Hamed, Jacob Bourjaily, Freddy Cachazo, Alexander Goncharov, Alexander Postnikov, Jaroslav Trnka, Grassmannian Geometry of Scattering Amplitudes (Cambridge University Press, 2016).
- [59] L. K. Williams, The positive Grassmannian, the amplituhedron, and cluster algebras, (2021), arXiv:2110.10856.
- [60] This fact can be proven by extracting the A_{∞} products via homological perturbation theory of [32, 38]. However, the products obtained this way are very complicated and do not immediately reveal neither the relation to convex geometry nor to pre-Calabi-Yau algebras. The details of the derivation will be given elsewhere [98].
- [61] E. Skvortsov and R. Van Dongen, Minimal models of field theories: SDYM and SDGR, (2022), arXiv:2204.09313 [hep-th].
- [62] Having well-defined vertices explicitly is very important. For example, [99] gives another type of homological perturbation theory in the same HiSGRA context, but the original recipe [99] to extract interactions leads to ill-defined vertices [100] (e.g. generic holographic correlation functions are infinite). From the field theory point of view [99] is not a concrete theory, but a general ansatz for interactions. It is not clear how to solve this problem since the duals of vector models are too nonlocal [8–12] to be treated by the standard tools. There are successful attempts to fix the first few local vertices (see [101, 102] and refs therein) which, however, do not address the nonlocal ones where the actual problem resides.
- [63] T. Tran, Toward a twistor action for chiral higher-spin gravity, (2022), arXiv:2209.00925 [hep-th].
- [64] E. Skvortsov and Y. Yin, On (spinor)-helicity and bosonization in AdS_4/CFT_3 , (2022), arXiv:2207.06976 [hep-th].
- [65] J. Maldacena and A. Zhiboedov, Constraining conformal field theories with a slightly broken higher spin symmetry, (2012), arXiv:1204.3882 [hep-th].
- [66] O. Aharony, G. Gur-Ari, and R. Yacoby, Correlation Functions of Large N Chern-Simons-Matter Theories and Bosonization in Three Dimensions, JHEP 12, 028, arXiv:1207.4593 [hep-th].
- [67] O. Aharony, Baryons, monopoles and dualities in Chern-Simons-matter theories, JHEP 02, 093, arXiv:1512.00161 [hep-th].
- [68] A. Karch and D. Tong, Particle-Vortex Duality from 3d Bosonization, Phys. Rev. X6, 031043 (2016), arXiv:1606.01893 [hep-th].
- [69] N. Seiberg, T. Senthil, C. Wang, and E. Witten, A Duality Web in 2+1 Dimensions and Condensed Matter Physics, Annals Phys. 374, 395 (2016), arXiv:1606.01989 [hep-th].

- [70] E. Sezgin and P. Sundell, Supersymmetric Higher Spin Theories, (2012), arXiv:1208.6019 [hep-th].
- [71] A. A. Sharapov and E. D. Skvortsov, Formal higher-spin theories and Kontsevich-Shoikhet-Tsygan formal-ity, Nucl. Phys. B921, 538 (2017), arXiv:1702.08218 [hep-th].
- [72] A. Sharapov and E. Skvortsov, Integrable Models From Non-Commutative Geometry With Applications to 3D Dualities, in 21st Hellenic School and Workshops on Elementary Particle Physics and Gravity (2022) arXiv:2204.08903 [hep-th].
- [73] A. S. Cattaneo and G. Felder, A Path integral approach to the Kontsevich quantization formula, Commun. Math. Phys. 212, 591 (2000), arXiv:math/9902090.
- [74] M. Blencowe, A Consistent Interacting Massless Higher Spin Field Theory in D=(2+1), Class.Quant.Grav. **6**, 443 (1989).
- [75] E. Bergshoeff, M. P. Blencowe, and K. S. Stelle, Area Preserving Diffeomorphisms and Higher Spin Algebra, Commun. Math. Phys. 128, 213 (1990).
- [76] A. Campoleoni, S. Fredenhagen, S. Pfenninger, and S. Theisen, Asymptotic symmetries of threedimensional gravity coupled to higher-spin fields, JHEP 1011, 007, arXiv:1008.4744 [hep-th].
- [77] M. Henneaux and S.-J. Rey, Nonlinear W_{∞} as Asymptotic Symmetry of Three-Dimensional Higher Spin Anti-de Sitter Gravity, JHEP **1012**, 007, arXiv:1008.4579 [hep-th].
- [78] C. N. Pope and P. K. Townsend, Conformal Higher Spin in (2+1)-dimensions, Phys. Lett. B225, 245 (1989).
- [79] E. S. Fradkin and V. Ya. Linetsky, A Superconformal Theory of Massless Higher Spin Fields in D = (2+1), Mod. Phys. Lett. **A4**, 731 (1989), [Annals Phys.198,293(1990)].
- [80] M. Grigoriev, I. Lovrekovic, and E. Skvortsov, New Conformal Higher Spin Gravities in 3d, JHEP 01, 059, arXiv:1909.13305 [hep-th].
- [81] A. Y. Segal, Conformal higher spin theory, Nucl. Phys. B664, 59 (2003), arXiv:hep-th/0207212 [hep-th].
- [82] A. A. Tseytlin, On limits of superstring in $AdS_5 \times S^5$, Theor. Math. Phys. **133**, 1376 (2002), [Teor. Mat. Fiz.133,69(2002)], arXiv:hep-th/0201112 [hep-th].
- [83] X. Bekaert, E. Joung, and J. Mourad, Effective action in a higher-spin background, JHEP 02, 048, arXiv:1012.2103 [hep-th].
- [84] P. Hähnel and T. McLoughlin, Conformal higher spin theory and twistor space actions, J. Phys. A 50, 485401 (2017), arXiv:1604.08209 [hep-th].
- [85] T. Adamo, P. Hähnel, and T. McLoughlin, Conformal higher spin scattering amplitudes from twistor space, JHEP 04, 021, arXiv:1611.06200 [hep-th].
- [86] T. Tran, Twistor constructions for higher-spin extensions of (self-dual) Yang-Mills, JHEP 11, 117, arXiv:2107.04500 [hep-th].
- [87] M. Sperling and H. C. Steinacker, Covariant 4dimensional fuzzy spheres, matrix models and higher spin, J. Phys. A50, 375202 (2017), arXiv:1704.02863 [hep-th].
- [88] H. Steinacker and T. Tran, A Twistorial Description of the IKKT-Matrix Model, (2022), arXiv:2203.05436 [hep-th].
- [89] R. de Mello Koch, A. Jevicki, K. Suzuki, and J. Yoon, AdS Maps and Diagrams of Bi-local Holography, JHEP 03, 133, arXiv:1810.02332 [hep-th].

- [90] O. Aharony, S. M. Chester, and E. Y. Urbach, A Derivation of AdS/CFT for Vector Models, (2020), arXiv:2011.06328 [hep-th].
- [91] D. Sullivan, Infinitesimal computations in topology, Publ. Math. IHES 47, 269 (1977).
- [92] P. van Nieuwenhuizen, Free graded differential superalgebras, in Group Theoretical Methods in Physics. Proceedings, 11th International Colloquium, Istanbul, Turkey, August 23-28, 1982 (1982) pp. 228-247.
- [93] R. D'Auria, P. Fre, and T. Regge, Graded Lie Algebra Cohomology and Supergravity, Riv. Nuovo Cim. 3N12, 1 (1980).
- [94] M. A. Vasiliev, Consistent equations for interacting massless fields of all spins in the first order in curvatures, Annals Phys. 190, 59 (1989).
- [95] M. Alexandrov, M. Kontsevich, A. Schwarz, and O. Zaboronsky, The Geometry of the Master Equation and Topological Quantum Field Theory, Int. J. Mod. Phys. A12, 1405 (1997), arXiv:hep-th/9502010 [hep-th].
- [96] G. Barnich and M. Grigoriev, BRST extension of the non-linear unfolded formalism, in *International School* / Seminar on Quantum Field Theory, Supersymmetry,

- High Spin Fields, Gravity Tomsk, Russia, March 20-26, 2005 (2005) arXiv:hep-th/0504119 [hep-th].
- [97] M. A. Vasiliev, Free massless fields of arbitrary spin in the de sitter space and initial data for a higher spin superalgebra, Fortsch. Phys. 35, 741 (1987).
- [98] A. Sharapov, E. Skvortsov, A. Sukhanov, and R. Van Dongen, More on Chiral Higher Spin Gravity and Convex Geometry, (2022), arXiv:2209.15441 [hepth].
- [99] M. A. Vasiliev, Closed equations for interacting gauge fields of all spins, JETP Lett. 51, 503 (1990).
- [100] N. Boulanger, P. Kessel, E. D. Skvortsov, and M. Taronna, Higher spin interactions in fourdimensions: Vasiliev versus Fronsdal, J. Phys. A49, 095402 (2016), arXiv:1508.04139 [hep-th].
- [101] V. E. Didenko, O. A. Gelfond, A. V. Korybut, and M. A. Vasiliev, Spin-locality of η^2 and $\overline{\eta}^2$ quartic higher-spin vertices, JHEP **12**, 184, arXiv:2009.02811 [hep-th].
- [102] O. A. Gelfond and A. V. Korybut, Manifest form of the spin-local higher-spin vertex $\Upsilon^{\eta \eta}_{\omega CCC}$, Eur. Phys. J. C 81, 605 (2021), arXiv:2101.01683 [hep-th].