

# The time has come for revising the rules of clozapine blood monitoring in Europe. A joint expert statement from the European Clozapine Task Force

*Short title: clozapine blood monitoring rules in Europe*

Hélène VERDOUX<sup>1</sup>, Robert A. BITTNER<sup>2</sup>, Alkomiet HASAN<sup>3,4</sup>, Mishal QUBAD<sup>2</sup>, Elias WAGNER<sup>3,5</sup>, Alexis LEPETIT<sup>6</sup>, Manuel ARROJO-ROMERO<sup>7</sup>, Christian BACHMANN<sup>8</sup>, Marieke BEEEX-OOSTERHUIS<sup>9</sup>, Jan BOGERS<sup>10</sup>, Andreja CELOFIGA<sup>11</sup>, Dan COHEN<sup>12</sup>, Domenico DE BERARDIS<sup>13</sup>, Marc DE HERT<sup>14</sup>, Carlos DE LAS CUEVAS<sup>15</sup>, Bjørn H. EBDROP<sup>16</sup>, Konstantinos N. FOUNTOULAKIS<sup>17</sup>, Daniel GUINART<sup>18</sup>, Dolores KEATING<sup>19</sup>, Miloslav KOPEČEK<sup>20</sup>, John LALLY<sup>21</sup>, Judit LAZÁRY<sup>22</sup>, Jurjen J. LUYKX<sup>23</sup>, Olalla MARONAS AMIGO<sup>24</sup>, Espen MOLDEN<sup>25</sup>, Jimmi NIELSEN<sup>26</sup>, Brian O'DONOGHUE<sup>27</sup>, Pierre OSWALD<sup>28</sup>, Flavian S. RADULESCU<sup>29</sup>, Christopher ROHDE<sup>30</sup>, Marina SAGUD<sup>31</sup>, Emilio J. SANZ<sup>32</sup>, Ivona ŠIMUNOVIĆ FILIPČIĆ<sup>33</sup>, Iris E. SOMMER<sup>34</sup>, Heidi TAIPALE<sup>35,36,37</sup>, Jari TIIHONEN<sup>35,36,38</sup>, Heli TUPPURAINEN<sup>35</sup>, Selene VEERMAN<sup>39</sup>, Alina WILKOWSKA<sup>40</sup>, Edoardo SPINA<sup>41</sup>, Peter SCHULTE<sup>42</sup>

1. French Clozapine Task Force; Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, team pharmacoepidemiology, UMR 1219, F-33000 Bordeaux, France

2. Goethe University Frankfurt, University Hospital, Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, Germany

3. Department of Psychiatry, Psychotherapy and Psychosomatics, Medical Faculty, University of Augsburg, Augsburg, Germany.

4. DZPG (German Center for Mental Health), partner site Munich/Augsburg, Germany.

5. Evidence-based Psychiatry and Psychotherapy, Faculty of Medicine, University of Augsburg, Augsburg, Germany.

6. French Clozapine Task Force; ACPA group, F-69340 Francheville; Hospices Civils de Lyon, Hôpital des Charpennes, F-69100 Villeurbanne; Institut des Sciences Cognitives Marc Jeannerod, CNRS, Team Neuroplasticity in Parkinson's disease, UMR 5229, F-69500 Bron, France

7. Department of Psychiatry, Complejo Hospitalario Universitario de Santiago, Santiago de Compostela, Spain

8. Department of Child and Adolescent Psychiatry, Ulm University, Germany

9. Department of Hospital Pharmacy, Albert Schweitzer Hospital, Dordrecht, The Netherlands

10. Mental health Organization Rivierduinen, Leiden, The Netherlands

11. Department of Psychiatry, University Medical Centre Maribor, Maribor, Slovenia

12. Dutch Clozapine Collaboration Group, Alkmaar, The Netherlands; Esdege-Reigersdaal, Heerhugowaard, The Netherlands

13. NHS, Department of Mental Health, G. Mazzini Hospital, Teramo, Italy

14. University Psychiatric Center Katholieke Universiteit Leuven, Leuvensesteenweg 517, 3070, Kortenberg, Belgium; Department of Neurosciences, Centre for Clinical Psychiatry, Katholieke Universiteit Leuven, Leuven, Belgium; Leuven Brain Institute, Katholieke Universiteit Leuven, Leuven, Belgium; Antwerp Health Law and Ethics Chair, University of Antwerp, Antwerp, Belgium.

15. Department of Internal Medicine, Dermatology and Psychiatry, School of Medicine, University of La Laguna, Canary Islands, Spain

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

16. Center for Neuropsychiatric Schizophrenia Research (CNSR), Mental Health Center, Glostrup, Copenhagen University Hospital – Mental Health Services CPH, Copenhagen, Denmark; Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark
17. 3<sup>rd</sup> Department of Psychiatry, Division of Neurosciences, School of Medicine, Faculty of Health Sciences, Aristotle University of Thessaloniki, Greece
18. Institut de Salut Mental, Hospital del Mar, Barcelona, Spain; Programa Recerca en Salut Mental, Hospital del Mar Research Institute, CIBERSAM, Barcelona; The Donald and Barbara Zucker School of Medicine, Northwell Health, New York, USA
19. Saint John of God Hospital, Stillorgan, Co. Dublin, Ireland
20. National Institute of Mental Health, Klecany, Czech Republic ;Department of Psychiatry, Charles University, Third Faculty of Medicine, Prague, Czech Republic
21. Department of Psychiatry, School of Medicine and Medical Science, University College Dublin, Dublin, Ireland; Department of Psychiatry, St Vincent's Hospital Fairview, Dublin, Ireland; Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience (IoPPN), King's College London, London, UK
22. Nyíró Gyula National Institute of Psychiatry and Addictions, Budapest, Hungary
23. Department of Psychiatry, Amsterdam UMC, Amsterdam, The Netherlands. GGZ inGeest Mental Health Care, Amsterdam, The Netherlands; Amsterdam Neuroscience (Mood, Anxiety, Psychosis, Stress & Sleep program) and Amsterdam Public Health (Mental Health program) research institutes, Amsterdam, The Netherlands; Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University Medical Center, Maastricht, the Netherlands
24. Pharmacogenomics and drug discovery (GenDeM), Health Research Institute of Santiago de Compostela (IDIS), Santiago de Compostela, Spain; Genomic Medicine Group, CIMUS, University of Santiago de Compostela, Santiago de Compostela, Spain; Centre for Biomedical Network Research on Rare Diseases (CIBERER), Instituto de Salud Carlos III, Madrid, Spain; Galician Public Foundation of Genomic Medicine (FPGMX), Galician Healthcare Service (SERGAS), Santiago de Compostela, Spain
25. Section for Pharmacology and Pharmaceutical Biosciences, Department of Pharmacy, University of Oslo, Oslo, Norway; Center for Psychopharmacology, Diakonhjemmet Hospital, Oslo, Norway.
26. Mental Health Centre Glostrup, Copenhagen University Hospital, Copenhagen, Denmark
27. Department of Psychiatry, University College Dublin, Dublin; Department of Psychiatry, St Vincents University Hospital, Dublin, Ireland
28. Department of psychiatry, Hôpital Universitaire de Bruxelles, Belgium
29. Center for Drug Sciences, Faculty of Pharmacy, University of Medicine and Pharmacy Carol Davila, Bucharest, Romania
30. Department of Affective Disorders, Aarhus University Hospital - Psychiatry, Aarhus, Denmark, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark
31. Department of Psychiatry, School of Medicine, University of Zagreb, Zagreb, Croatia; Department for Psychiatry and Psychological Medicine, University Hospital Center Zagreb, Croatia
32. Department of Physical Medicine and Pharmacology, School of Medicine, Universidad de La Laguna, Canary Islands, Spain; Hospital Universitario de Canarias, Tenerife, Spain
33. Department of Psychiatry and Psychological Medicine, University Hospital Centre Zagreb; Faculty of Dental Medicine and Health, Josip Juraj Strossmayer University of Osijek, Osijek, Croatia
34. Department of Neuroscience, and Department of Psychiatry, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands
35. Department of Forensic Psychiatry, University of Eastern Finland, Niuvanniemi Hospital, 70240 Kuopio, Finland
36. Department of Clinical Neuroscience, Karolinska Institutet, and Center for Psychiatry Research, Stockholm City Council, Stockholm, Sweden
37. School of Pharmacy, University of Eastern Finland, Kuopio, Finland
38. Neuroscience Center, University of Helsinki, Helsinki, Finland
39. Mental Health Services Noord-Holland-Noord, Schagen, the Netherlands; Dutch Clozapine Collaboration Group, Alkmaar, the Netherlands
40. Department of Psychiatry, Medical University of Gdańsk, Gdańsk, Poland
41. Department of Clinical and Experimental Medicine, University of Messina, Messina, Italy
42. Mental Health Services Noord-Holland-Noord, Alkmaar, the Netherlands; Dutch Clozapine Collaboration Group, Alkmaar, the Netherlands

1  
2  
3  
4  
5  
6

**Correspondence:** Prof Hélène Verdoux, University Bordeaux, UMR 1219, Case 36, 146 Rue Léo  
Saignat, 33076, Bordeaux Cedex  
helene.verdoux@u-bordeaux.fr

## Abstract

The European Clozapine Task Force is a group of psychiatrists and pharmacologists practicing in 18 countries under European Medicines Agency (EMA) regulation, who are deeply concerned about the underuse of clozapine in European countries. Although clozapine is the most effective antipsychotic for people with treatment-resistant schizophrenia, a large proportion of them do not have access to this treatment. Concerns about clozapine-induced agranulocytosis and stringent blood monitoring rules are major barriers to clozapine prescribing and use. There is a growing body of evidence that the incidence of clozapine-induced agranulocytosis is very low after the first year of treatment. Maintaining lifelong monthly blood monitoring after this period contributes to unjustified discontinuation of clozapine. We leverage recent and replicated evidence on the long-term safety of clozapine to call for the revision and updating of the EMA's blood monitoring rules, thus aiming to overcome this major barrier to clozapine prescribing and use. We believe the time has come for relaxing the rules without increasing the risks for people using clozapine in Europe.

**Key Words:** Clozapine, treatment-resistant schizophrenia, blood monitoring, regulation

## 1      **Introduction**

2            Shortly after the introduction of clozapine in the seventies, several cases of fatal  
3      agranulocytosis in Finland led to its market withdrawal in almost all countries. Following the  
4      study by Kane and colleagues in people with treatment-resistant schizophrenia (TRS) [1],  
5      clozapine was reintroduced with strict blood monitoring rules to detect incident  
6      agranulocytosis, estimated at 1-2% lifetime. More than 35 years later, these rules have not  
7      been revised in most countries [2]. Under current European Medicines Agency (EMA)  
8      regulation, white blood cell count (WBC) and absolute neutrophil count (ANC) monitoring is  
9      required weekly for the first 18 weeks after initiation of treatment and then monthly for the  
10     duration of treatment. For many patients this can mean decades of monthly blood sampling.

11           Clozapine remains the only approved antipsychotic for TRS with superior efficacy and  
12     effectiveness for several endpoints, such as suicide, psychotic symptoms, relapse,  
13     rehospitalization, medication adherence, aggression or substance use. Clozapine use is also  
14     associated with reduced all-cause, suicide and cardiovascular mortality. While TRS occurs in  
15     about one third of schizophrenia patients, only a minority of them are prescribed this  
16     medication [3], which represents a missed opportunity for these people, as their chances of  
17     recovery are much less favorable without clozapine [3-7].

18           The need of continued blood tests “for life” is a significant barrier to clozapine  
19     initiation and maintenance treatment increasing the risk of premature discontinuation [8,9].  
20     From the prescriber's perspective, the most common barriers to clozapine initiation are  
21     related to the institutional complexity of mandatory blood monitoring and an overestimated  
22     users' poor adherence to this monitoring [9]. Prescribers' overestimation of agranulocytosis

1 risk also contribute to under-monitoring of other adverse drug reactions with higher lethality  
2 including myocarditis, pneumonia or ileus [10].

3 Blood monitoring also comes at a cost. In addition to the direct costs of blood tests,  
4 monitoring is time-consuming for laboratories and pharmacies. It requires complex  
5 organizational adaptations to guarantee continuity of care, particularly at a time when  
6 medical and paramedical staffing levels are low. A cost-effectiveness analysis compared  
7 several blood monitoring strategies in a theoretical cohort of 100,000 people treated with  
8 clozapine [10]. While the « no monitoring » strategy was the most cost-effective, no  
9 difference was found regarding quality-adjusted life-year gained compared to the other  
10 modalities, irrespective of the stringency of blood monitoring rules.

#### 11 **Low risk of agranulocytosis after first year of treatment**

12 Recent studies have confirmed that the risk window for clozapine-related  
13 agranulocytosis is mostly limited to the first months of treatment. A meta-analysis of 108  
14 studies (1983-2017, 448,647 clozapine users) found a 3.8% incidence of mild neutropenia ( $\leq$   
15  $1.5 \times 10^9$  per L), a 1.3% incidence of moderate neutropenia ( $\leq 1.0 \times 10^9$  per L) and a 0.9% of  
16 severe neutropenia, commonly referred to as agranulocytosis ( $\leq 0.5 \times 10^9$  per L) (0.7% when  
17 only high-quality study were considered) [5]. The agranulocytosis fatality rate was one death  
18 for 7,700 clozapine users. The agranulocytosis prevalence was identical in studies conducted  
19 before and after the introduction of blood monitoring in 1990. In most cases (75%), mild  
20 neutropenia did not progress to severe neutropenia. Most cases of agranulocytosis occurred  
21 in the first few months of treatment: 38% in the first month, 56% within two months, 84%  
22 within the first 18 weeks and 89% within the first year.

Neutropenia  $<1.0 \times 10^9$  per L occurred in 1.2% of users in a population-based study of 26,000 clozapine users in Australia and New Zealand (1990-2022) with not fatal cases [6]. The risk was highest during the first 18 weeks of treatment (weekly incidence 0.13%) and became negligible after 24 months. The incidence of any neutropenic event was very low when clozapine was reintroduced in people with no history of neutropenia over two years of cumulative monitoring. This finding suggests that there is no need to resume a weekly monitoring schedule after a clozapine interruption in people with no history of neutropenia.

Another population-based study of 14,037 clozapine users in Finland (1996-2017) found a cumulative incidence of agranulocytosis of 1.37 % with clozapine compared with 0.13% with other antipsychotics [7]. The mortality rate for agranulocytosis was one death in 3,559 people starting clozapine. Compared to modal treatment duration for non-clozapine antipsychotics (12–23 months), the risk of agranulocytosis decreased over time from adjusted odds ratio of 36 during the first six months of clozapine exposure to 4.38 for exposure  $\geq 54$  months: it became then comparable to that observed over the first six months of treatment for other antipsychotics for which no mandatory monitoring is required.

### **Stringency of blood monitoring rules already differs between European countries**

A review of clozapine monitoring regulation in 102 countries highlighted the wide variability in monitoring rules and in WBC/ANC criteria for stopping clozapine [2]. Although EMA regulation applies to all EU countries as well as Norway and Iceland, the blood monitoring rules already differ from one country to another [2] (supplementary Table 1). In several countries, the rule "no blood, no drug" does not apply and monitoring is already relaxed. The recommendations of the Netherlands Clozapine Collaboration Group, set up in 2006, allow for off-label less stringent monitoring rules after the first 18 weeks of treatment

1 if the prescriber and the well-informed patient decide so. Similar recommendations apply to  
2 Iceland with the possibility of relaxed monitoring after six months. Monitoring stringency has  
3 no impact on the incidence of agranulocytosis but is inversely associated with the rate of  
4 clozapine use in each country [2].

5 Criteria for clozapine discontinuation following a drop in WBC count also vary from  
6 country to country. Too stringent criteria may lead to unjustified discontinuation due to  
7 transient neutropenia not related to clozapine. Interdisciplinary boards involving psychiatrists  
8 and hematologists would be needed to exclude concurrent reasons for ANC decrease.

9 In most European countries, mild neutropenia i.e.  $ANC < 1.5 \times 10^9$  per L requires  
10 discontinuation (supplementary Table 1). However, in a few European countries under EMA  
11 regulation, the US Food and Drug Administration (FDA) criteria ( $ANC < 1.0 \times 10^9$  per L) are used  
12 after 18 weeks [2].

13 Several countries have adjusted the threshold criteria for clozapine discontinuation  
14 for people concerned with benign ethnic neutropenia (BEN) [1] since their access to clozapine  
15 is markedly restricted under the standard criteria. For instance, the FDA threshold criteria for  
16 clozapine discontinuation for BEN is  $ANC < 0.5 \times 10^9$  per L. The BEN adjusted criteria are not  
17 yet applied in any country under the EMA regulation, leading to discrimination regarding  
18 access to clozapine for the many people living with BEN in Europe. For example, up to 80% of  
19 people black African or Caribbean descent have the Duffy-null phenotype of the ACKR1  
20 (Atypical Chemokine Receptor 1) gene associated with BEN.

### 21 **Reducing barriers to clozapine by revising European blood monitoring rules**

22 35 years ago, stringent blood monitoring rules were justified to ensure the successful  
23 reintroduction of clozapine. Today, this stringency no longer matches the reality of the actual



1 hematological risk, as demonstrated by the growing body of evidence on the temporal  
2 pattern of clozapine-induced agranulocytosis. No other drug approved in Europe with a  
3 negligible risk of agranulocytosis after the first year of treatment requires lifelong monthly  
4 blood monitoring.

5 The European Clozapine Task Force is a group of psychiatrists and pharmacologists  
6 practicing in 18 countries under EMA regulation with extensive clinical and research expertise  
7 in clozapine. We are deeply concerned about the underuse of clozapine in European countries.  
8 We are convinced revising and updating of the EMA's blood monitoring rules could help to  
9 overcome the major barrier to the use of clozapine [9].

10 Our proposals for new clozapine blood monitoring rules are detailed in Table 1. Only  
11 ANC criteria are given as the majority of authors (60%) are in favour of restricting mandatory  
12 monitoring to ANC based on FDA regulation revisions in 2015. However, no consensus could  
13 be reached among the members of the European Clozapine Task Force on this point.

14 Based on the currently available evidence, we propose to:

- 15 (i) relax the blood monitoring schedule after 12 months and 24 months of treatment,
- 16 (ii) lower ANC threshold for clozapine initiation and discontinuation,
- 17 (iii) use an adjusted threshold for BEN,
- 18 (iv) relax monitoring schedule after clozapine interruption,
- 19 (v) harmonize (i) to (iv) across Europe.

20 The benefits of revising the monitoring rules outweigh the potential risks associated  
21 with less stringent rules: allowing more people with TRS to get clozapine will save lives  
22 without increasing the risk of agranulocytosis.

## Conclusion

Access to clozapine is currently severely hampered by multiple barriers resulting in a real loss of opportunity for many people with TRS, with a high price being paid by both the users and the healthcare system. Whenever deemed indicated, clozapine treatment must be initiated as soon as possible to promote recovery and to increase life expectancy. Better addressing these unmet needs should be considered as a public health priority by European health regulatory agencies, as it is currently done by the FDA. More balanced monitoring rules would contribute to reduce clozapine underprescription as well as discrimination against people with BEN. The time has come for revising the rules of clozapine blood monitoring in Europe.

**Financial Support.** This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

**Acknowledgements.** Not relevant.

**Competing interests.** The following authors declare none: Hélène VERDOUX, Robert BITTNER, Alexis LEPETIT, Manuel ARROJO-ROMERO, Christian BACHMANN, Marieke BEEX-OOSTERHUIS, Jan BOGERS, Dan COHEN, Marc DE HERT, Carlos DE LAS CUEVAS, John LALLY, Dolores KEATING, Judit LAZÁRY, Jurjen J. LUYKX, Olalla MARONAS AMIGO, Jimmi NIELSEN, Brian O'DONOGHUE, Pierre OSWALD, Flavian S RADULESCU, Christopher ROHDE, Emilio J. SANZ, Ivona ŠIMUNOVIĆ FILIPČIĆ, Iris E. SOMMER, Selene VEERMAN, Edoardo SPINA, Peter SCHULTE; Alkomiet HASAN is editor of the German (DGPPN) schizophrenia treatment guidelines, first author of the WFSBP schizophrenia treatment guidelines; on advisory boards of and speaker fees from AbbVie, Advanz (speaker fees only), Janssen-Cilag, Lundbeck, Recordati, Rovi, Teva, and Otsuka; Mishal QUBAD has received speaker fees from Recordati; Elias WAGNER was invited to advisory boards from Recordati, Teva and Boehringer Ingelheim; Andreja CELOFIGA participated in lectures for the following companies: Krka, Bonifar, Viatrix, Gedeon Richter, Lek, and Lundbeck-Pharma; Domenico DE BERARDIS was speaker/consultant for Angelini, Innova Pharma, Janssen, Lundbeck, Viatrix, Rovi and Otsuka Konstantinos; Björn EBDROP is part of the Advisory Board of Boehringer Ingelheim, Lundbeck Pharma A/S; and has received lecture fees from Boehringer Ingelheim, Otsuka Pharma Scandinavia AB, and Lundbeck Pharma A/S; N. FOUNTOLAKIS has received in the past grants, served as a consultant, advisor, or CME speaker, or received support to attend congresses by the following entities: AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Ferrer, Gedeon Richter, Janssen, Lundbeck, Otsuka, Pfizer, the Pfizer Foundation, Sanofi-Aventis, Servier, Shire, and others; Daniel GUINART has been a consultant and/or advisor of has received honoraria from Angelini, Janssen, Lundbeck/Lundbeck Foundation, Otsuka, Teva ; he currently receives funding from the European Commission (HORIZON EUROPE) and from the Instituto de Salud Carlos III (ISCIII) ; Miloslav KOPEČEK is member of advisory board and speakers' honoraria from Angelini Pharma and Lundbeck; Espen MOLDEN has received speaker's honorarium from Lundbeck and Otsuka; Marina SAGUD has presented lectures for the following companies: Angelini, Belupo, Gedeon Richter, Johnson & Johnson, Krka Farma d.o.o., Lundbeck, Pliva d.o.o., and Viatrix; participated in clinical trials as a principal investigator for Boehringer Ingelheim; and is a member of the advisory board of Lundbeck and Pliva; Heidi TAIPALE has participated in research projects funded by grants from Janssen to her employing institution; and has received

1 lecture fees from Gedeon Richter, Janssen, Lundbeck and Otsuka; Jari TIIHONEN has participated in research  
2 projects funded by grants from Janssen to his employing institution; has served as a consultant for Healthcare  
3 Global Village, HLS Therapeutics, Janssen, Lundbeck, Orion, Teva and WebMD Global; and has received  
4 honoraria from Janssen, Lundbeck, and Otsuka; Heli TUPPURAINEN has received speakers fee from Lundbeck ;  
5 Alina WILKOWSKA has received research support from Angelini, Biogen, Eli Lilly and Company, Janssen- Cilag,  
6 Lundbeck, Polpharma, Sanofi, Termedia and Valeant.

7  
8 **Data availability statement.** Not relevant.  
9  
10

## References

1. Kane J, Honigfeld G, Singer J, et al. Clozapine for the treatment-resistant schizophrenic. A double-blind comparison with chlorpromazine. *Arch Gen Psychiatry*. 1988;45:789-96. 10.1001/archpsyc.1988.01800330013001
2. Oloyede E, Blackman G, Whiskey E, et al. Clozapine haematological monitoring for neutropenia: a global perspective. *Epidemiol Psychiatr Sci*. 2022;31:e83. 10.1017/s204579602200066x
3. Bachmann CJ, Aagaard L, Bernardo M, et al. International trends in clozapine use: a study in 17 countries. *Acta Psychiatr Scand*. 2017;136:37-51. 10.1111/acps.12742
4. Bittner RA, Reif A, Qubad M. The ever-growing case for clozapine in the treatment of schizophrenia: an obligation for psychiatrists and psychiatry. *Curr Opin Psychiatry*. 2023;36:327-336. 10.1097/ycp.0000000000000871
5. Myles N, Myles H, Xia S, et al. Meta-analysis examining the epidemiology of clozapine-associated neutropenia. *Acta Psychiatr Scand*. 2018;138:101-109. 10.1111/acps.12898
6. Northwood K, Myles N, Clark SR, et al. Evaluating the epidemiology of clozapine-associated neutropenia among people on clozapine across Australia and Aotearoa New Zealand: a retrospective cohort study. *Lancet Psychiatry*. 2024;11:27-35. 10.1016/s2215-0366(23)00343-7
7. Rubio JM, Kane JM, Tanskanen A, et al. Long-term persistence of the risk of agranulocytosis with clozapine compared with other antipsychotics: a nationwide cohort and case-control study in Finland. *Lancet Psychiatry*. 2024;11:443-450. 10.1016/s2215-0366(24)00097-x
8. Schulte PFJ, Veerman SRT, Bakker B, et al. Risk of clozapine-associated agranulocytosis and mandatory white blood cell monitoring: Can the regulations be relaxed? *Schizophr Res*. 2024;268:74-81. 10.1016/j.schres.2023.09.024
9. Verdoux H, Quiles C, Bachmann CJ, et al. Prescriber and institutional barriers and facilitators of clozapine use: A systematic review. *Schizophr Res*. 2018;201:10-19. 10.1016/j.schres.2018.05.046
10. Girardin FR, Poncet A, Blondon M, et al. Monitoring white blood cell count in adult patients with schizophrenia who are taking clozapine: a cost-effectiveness analysis. *Lancet Psychiatry*. 2014;1:55-62. 10.1016/s2215-0366(14)70245-7

**Table 1. European Clozapine Task Force proposals for new clozapine blood monitoring rules in countries under European Medicines Agency regulation**

Current SPC <sup>1</sup>		Proposals <sup>2</sup>
<b>Mandatory routine blood monitoring schedule</b>	<b>WBC<sup>3</sup> and ANC<sup>4</sup></b> <ul style="list-style-type: none"> <li>○ baseline before initiation</li> <li>○ weekly for 18 weeks after initiation</li> <li>○ then monthly irrespective of treatment duration</li> </ul>	<b>First 12 months</b> ANC <ul style="list-style-type: none"> <li>○ baseline before initiation</li> <li>○ weekly for 18 weeks after initiation</li> <li>○ then monthly for 34 weeks</li> </ul> <b>After 12 months<sup>5</sup></b> <ul style="list-style-type: none"> <li>○ ANC every 12 weeks if no history of leukopenia or neutropenia during the first year</li> </ul> <b>After 24 months<sup>5</sup></b> <ul style="list-style-type: none"> <li>○ yearly ANC if no history of leukopenia or neutropenia during two years</li> </ul>
<b>Standard thresholds for</b>		
Initiation/continuation	<ul style="list-style-type: none"> <li>○ <math>ANC \geq 2.0 \times 10^9/L</math></li> <li>○ <math>WBC \geq 3.5 \times 10^9/L</math></li> </ul>	○ $ANC \geq 1.5 \times 10^9 \text{ per } L^6$
Monitoring twice a week	<ul style="list-style-type: none"> <li>○ <math>ANC 1.5-2 \times 10^9/L</math></li> <li>○ <math>WBC 3.0-3.5 \times 10^9/L</math></li> </ul>	○ $ANC 1.0-1.5 \times 10^9/L^6$
Discontinuation (red)	<ul style="list-style-type: none"> <li>○ <math>ANC &lt; 1.5 \times 10^9/L</math></li> <li>○ <math>WBC &lt; 3.0 \times 10^9/L</math></li> </ul>	○ $ANC < 1.0 \times 10^9 \text{ per } L^6$
<b>BEN<sup>7</sup> adjusted thresholds for</b>		
Initiation/continuation		○ $ANC \geq 1.0 \times 10^9 \text{ per } L^6$
Monitoring twice a week		○ $ANC 0.5-1.0 \times 10^9 \text{ per } L^6$
Discontinuation		○ $ANC < 0.5 \times 10^9 \text{ per } L^6$
<b>Monitoring schedule after clozapine interruption</b>	interruption > 3 days and < 4 weeks <ul style="list-style-type: none"> <li>○ weekly for 6 weeks then monthly</li> </ul> <i>interruption</i> > 4 weeks <ul style="list-style-type: none"> <li>○ weekly for 18 weeks then monthly</li> </ul>	irrespective of the duration of interruption <ul style="list-style-type: none"> <li>○ no need to resume weekly schedule if no history of neutropenia during two cumulative years of monitoring</li> </ul>

1. Summaries of Product Characteristics ; 2. Only Absolute Neutrophil Count criteria are given as the majority of authors (60%) are in favour of restricting mandatory monitoring to ANC based on Food and Drug Administration regulation revisions in 2015; however, no consensus could be reached among the members of the European Clozapine Task Force on this point ; 3. White Blood Cells count ; 4. Absolute Neutrophil Count ; 5. Even if the frequency of routine mandatory monitoring is reduced, ANC must be performed immediately in the event of possible symptoms of infection (e.g. fever, sore throat, mouth/throat ulcers). Additional ANC may be considered after addition of valproic acid to clozapine, especially during the initiation period. 6. Food and Drug Administration criteria; 7. Benign Ethnic Neutropenia (haematology consultation may be needed to confirm the diagnosis).